

Lab/UFR-HEP0605/GNPHE/0605/VACBT/0605

Pure fermionic twistor like model & target space supersymmetry

R. Ahl Laamara, L.B Drissi, H. Jehjoui, E H Saidi*

1. Lab/UFR- Physique des Hautes Energies, Faculté des Sciences, Rabat, Morocco,

2. GNPHE, Groupement National de Physique des Hautes Energies,

Siège focal: Faculté des Sciences de Rabat, Morocco.

3. Virtual African Centre for Basic Science and Technology, VACBT,

Focal point, LabUFR-PHE, Rabat, Morocco.

February 1, 2008

Abstract

Using world line fermions $\Upsilon_{\pm}^m = \Upsilon_{\pm}^m(\tau)$ valued in vector representation of $SO(d, 4-d)$ with $d = 2, 3, 4$, we develop a pure fermionic analog of Penrose twistor construction. First, we show that Fermi antisymmetry requiring $(\Upsilon_{\pm}^m)^2 = 0$ can be solved by using twistor like variables. Then we study the corresponding dual twistor like field action and show that quantum spectrum exhibits naturally $4D$ $\mathcal{N} = 1$ target space supersymmetry. Higher spin world line field solutions of the constraint $(\Pi_s^m)^2 = 0$, $s \in \mathbb{Z}$ are also discussed.

Keywords: World line (world sheet) supersymmetry, Twistor gauge theory, pure fermionic twistor like model, 4D target space supersymmetry.

*h-saidi@fsr.ac.ma

1 Introduction

During the few last years, a lot of developments has been made on twistor approach for describing $4D \mathcal{N} = 4$ supersymmetric Yang-Mills and $10D$ superstring theories in complex projective superspace $\mathbb{CP}^{(3|4)}$ [1]-[5], see also [6]-[10] for earlier related works. An important interest has been given to the study of twistor supersymmetric gauge theory and stringy extension [11]-[15]. The key issue behind this revival of interest in Penrose model [16]-[18] is the proposal of [1] where it was shown that the supertwistor space $\mathbb{CP}^{(3|4)}$ is a Calabi-Yau supermanifold [19]-[28] capturing naturally special properties of gauge theory correlation functions; in particular planar amplitudes of $4D \mathcal{N} = 4$ supersymmetric $U(N)$ Yang-Mills theory, and exhibiting the good properties of a target space for topological string theory on Calabi-Yau threefolds [29]- [35].

Motivated by recent results in this matter and particular aspects of the so called two-time physics [36], we propose in this paper to set up the basis of a pure world line fermionic extension of standard Penrose model. Our construction is based on solving the nilpotency property $(\Upsilon_m)^2 = 0$ of the world line fermions $\Upsilon_m = \Upsilon_m(\tau)$ by using a Penrose like method. Recall that twistor approach of Penrose is based, amongst others, on solving the condition $(p_m)^2 = 0$ of a massless particle in 4D- space time. It happens that, besides the striking analogy between the solutions of $(p_m)^2 = 0$ and $(\Upsilon_m)^2 = 0$ and their underlying twistor world line field formulations, we find moreover that fermionic twistor like space is a graded superspace capturing four dimension target space supersymmetry in a wonderful way.

To derive our pure fermionic twistor like model, we shall mimick the standard twistor derivation of Penrose [16] as reformulated in [1]. The main steps of this construction can, roughly, be summarized as follows:

(a) The usual real world line free massless field $x_\alpha^\dot{\alpha}(\tau)$ of standard Penrose twistor model is now replaced by a free world line fermion $\Upsilon_\alpha^\dot{\alpha}(\tau)$. Both of these world line fields are 4-vectors of the real $\mathbb{R}^{(d,4-d)}$ space, $d = 2, 3, 4$, with isometry group $SO(d, 4-d)$ and diagonal metric η_{mn} with $(d, 4-d)$ signature.

(b) The role of the conjugate momentum $p_\alpha^\alpha = \left(\frac{\delta \mathcal{S}_{BT}}{\delta \partial_\tau x_\alpha^\alpha} \right)$ of the field x_α^α gets played by $\pi_\alpha^\alpha = \left(\frac{\delta \mathcal{S}_{FT}}{\delta \partial_\tau \Upsilon_\alpha^\alpha} \right)$, the fermionic conjugate momentum of Υ_α^α . In these relations, $\mathcal{S}_{BT} = \mathcal{S}[X_\alpha^\alpha, p_\alpha^\alpha]$ and $\mathcal{S}_{FT} = \mathcal{S}[\Upsilon_\alpha^\alpha, \pi_\alpha^\alpha]$ stand for the world line field actions of the bosonic x_α^α , p_α^α and the fermionic Υ_α^α , π_α^α fields respectively.

(c) The role of the zero mass relation $p^2 = 0$ of the free field bosonic theory is now played by the fermionic nilpotency property $\pi^2 = 0$ of the fermionic field π_α^α . Notice that in our fermionic analog, there is a subtlety to announce. Besides $\text{Tr}(\pi^2) = 0$, there are two more constraint eqs one has to deal with. They are given by $\text{Tr}(\Upsilon^2) = 0$ and $\text{Tr}(\pi\Upsilon) = 0$

and it turns out that they are altogether captured by a local $SP(2, \mathbb{R})$ gauge invariance to be discussed in section 4.

The next step is to solve the constraint eqs using Penrose method and derive the dual twistor field action describing the fermionic fields. After that we study the quantization of this model and shows how $4D$ target space supersymmetry follows.

The presentation of this paper is as follows: In section 2, we develop a little bit our motivations and give preliminary results. In section 3, we consider superspace representations of world sheet $\mathfrak{n} = (1, 1)$ superalgebra and world line reductions. This super- invariance allows to make an idea on the basis of our construction and possible supersymmetric extension. In section 4, we give details on our fermionic twistor like analog. First we comment the standard Penrose construction for world line bosons. Then we build the pure fermionic model. In section 5, we solve the constraint eqs and derive the quantum fermionic twistor like field action. Section 6 is devoted to conclusion and discussions.

2 Motivations and preliminary results

To make a general idea on our pure fermionic analogue of Penrose twistor construction and on the method we will use to handle it, let us start by recalling some useful features. We take also this opportunity to give preliminary results that will be developed later.

Consider a free world line bosonic field,

$$x^m(\tau) \sim x_{\alpha}^{\dot{\alpha}}(\tau), \quad m = 0, 1, 2, 3, \quad \alpha, \dot{\alpha} = 1, 2 \quad (2.1)$$

moving inside the real four dimension space $\mathbb{R}^{d, 4-d}$ with $d = 2, 3, 4$. The field action $\mathcal{S}_{BT}$ describing the dynamics of the massless particle reads as,

$$\mathcal{S}_{BT}[x, p, e] = \int d\tau (p_m \partial_{\tau} x^m - e p^2), \quad (2.2)$$

where p_m is the conjugate momentum of the world line field x^m and $e = e(\tau)$ is the gauge field capturing the constraint eq fixing the mass squared $p^2 = M^2$ of the particle as,

$$p^2 = \eta^{mn} p_m p_n = 0. \quad (2.3)$$

To handle this equation, we rewrite the 4- component energy momentum vector p_m like a hermitian 2×2 matrix,

$$p_m \sim (\sigma_m)_{\alpha}^{\dot{\alpha}} p_{\dot{\alpha}}^{\alpha} \quad (2.4)$$

Here $p_{\dot{\alpha}}^{\alpha}$ is a bi-spinor of the $SO(d, 4-d)$ orthogonal group and the σ_m 's are the usual Pauli matrices satisfying the Clifford algebra. The diagonal metric η_{mn} of $\mathbb{R}^{d, 4-d}$ can be

also expressed in terms of the product of the invariant spinor metrics $\epsilon_{\alpha\beta}$ and $\epsilon_{\dot{\alpha}\dot{\beta}}$. Using dotted and undotted spinor indices, we can re-express functions involving the 4-vector p_m in terms of 2×2 matrix $p_{\dot{\alpha}}^{\alpha}$. For instance, the quadratic invariant p^2 is mapped to $\det(p_{\dot{\alpha}}^{\alpha})$ which can be also rewritten as $\text{Tr}(p^2)$ with,

$$\begin{aligned}\text{Tr}(p^2) &= \epsilon_{\alpha\beta} p_{\dot{\alpha}}^{\alpha} p_{\dot{\beta}}^{\beta} \epsilon^{\dot{\alpha}\dot{\beta}} = p_{\dot{\alpha}}^{\alpha} \cdot p_{\dot{\alpha}}^{\dot{\alpha}} = p_{\dot{\alpha}}^{\dot{\alpha}} \cdot p_{\dot{\alpha}}^{\alpha}, \\ \epsilon_{\alpha\beta} \epsilon^{\beta\gamma} &= \delta_{\alpha}^{\gamma}, \quad \epsilon^{\dot{\alpha}\dot{\beta}} \epsilon_{\dot{\beta}\dot{\gamma}} = \delta_{\dot{\gamma}}^{\dot{\alpha}}.\end{aligned}\tag{2.5}$$

From classical view, $p_m = p_m(\tau)$ is a world line field and the equation $\text{Tr}(p^2) = 0$ admits a remarkable solution given by,

$$p_{\dot{\alpha}}^{\alpha} = \lambda^{\alpha} \tilde{\lambda}_{\dot{\alpha}}, \quad \alpha, \dot{\alpha} = 1, 2, \tag{2.6}$$

where the world line fields $\lambda^{\alpha} = \lambda^{\alpha}(\tau)$ and $\tilde{\lambda}_{\dot{\alpha}} = \tilde{\lambda}_{\dot{\alpha}}(\tau)$ are $SO(d, 4-d)$ spinors with positive and negative chirality respectively.

$$\lambda^{\alpha} \sim \left(\frac{1}{2}, 0\right), \quad \tilde{\lambda}_{\dot{\alpha}} \sim \left(0, \frac{1}{2}\right). \tag{2.7}$$

By computing the invariant square $\text{Tr}(p^2)$ of the solution $p_{\dot{\alpha}}^{\alpha} = \lambda^{\alpha} \tilde{\lambda}_{\dot{\alpha}}$, we see that the vanishing condition,

$$\text{Tr}(p^2) = \lambda^2 \tilde{\lambda}^2 = 0, \tag{2.8}$$

is ensured by requiring

$$\lambda^2 = \epsilon_{\alpha\beta} \lambda^{\alpha} \lambda^{\beta}, \quad \tilde{\lambda}^2 = \epsilon^{\dot{\alpha}\dot{\beta}} \tilde{\lambda}_{\dot{\alpha}} \tilde{\lambda}_{\dot{\beta}} \tag{2.9}$$

to vanish identically; i.e

$$\lambda^2 = 0 \quad \text{and/or} \quad \tilde{\lambda}^2 = 0. \tag{2.10}$$

This requirement is naturally fulfilled in twistor analysis. Notice that the vanishing property of the invariant squares of λ^{α} and $\tilde{\lambda}_{\dot{\alpha}}$ ($\lambda^2 = 0$ and $\tilde{\lambda}^2 = 0$) is due to the antisymmetry property of the dotted and undotted metrics $\epsilon^{\dot{\alpha}\dot{\beta}} = -\epsilon^{\dot{\beta}\dot{\alpha}}$, $\epsilon_{\alpha\beta} = -\epsilon_{\beta\alpha}$ and because of the commutativity of the Penrose variables [16, 18]-[1, 27, 28],

$$\lambda^{\alpha} \lambda^{\beta} = \lambda^{\beta} \lambda^{\alpha}, \quad \text{and} \quad \tilde{\lambda}_{\dot{\alpha}} \tilde{\lambda}_{\dot{\beta}} = \tilde{\lambda}_{\dot{\beta}} \tilde{\lambda}_{\dot{\alpha}}. \tag{2.11}$$

However in looking carefully at this solution, one notes a set of remarkable properties indicating that Penrose representation for $p_{\dot{\alpha}}^{\alpha} = \lambda^{\alpha} \tilde{\lambda}_{\dot{\alpha}}$ is just a part of a general picture which should deserve more attention. Below we give three of them; they will be developed in much more details in forthcoming sections.

2.1 $p_{\dot{\alpha}}^{\alpha}$: First member of a field family $\{\Pi_{s\dot{\alpha}}^{\alpha}, s \in \mathbb{Z}\}$

First note that to solve the condition,

$$\lambda^2 \tilde{\lambda}^2 = 0, \quad (2.12)$$

it is, a priori, enough to require $\lambda^2 = 0$ whatever $\tilde{\lambda}^2$ is or inversely. But this is not exactly what happens since we have both $\lambda^2 = 0$ and $\tilde{\lambda}^2 = 0$. A careful study shows that this degeneracy hides in fact a remarkable property which can be used to go beyond the standard Penrose twistor construction for bosons. In field theory, eq(2.6) can be viewed as just the leading relation of a family of "conformal spin" s world line field variables. A class of these fields is given by the two following infinite sets:

$$(\Pi'_s)_{\dot{\alpha}}^{\alpha} = \lambda^{\alpha} \left(\tilde{\lambda}_s \right)_{\dot{\alpha}}, \quad s = 0, \pm 2, \pm 4, \dots, \quad (2.13)$$

for world line bosons and by

$$(\Pi'_s)_{\dot{\alpha}}^{\alpha} = \lambda^{\alpha} \left(\tilde{\zeta}_s \right)_{\dot{\alpha}}, \quad s = \pm 1, \pm 3, \pm 5, \dots, \quad (2.14)$$

for world line fermions. In these relations λ^{α} is as before; but $\tilde{\lambda}_{s\dot{\alpha}}$ and $\tilde{\zeta}_{s\dot{\alpha}}$ are new fields with even and odd conformal spins $\frac{s}{2}$ respectively. As a first result we want to mention here is that, whatever the value of s , all these $(\Pi'_s)_{\dot{\alpha}}^{\alpha}$, $s \in \mathbb{Z}$, exhibit the property,

$$\text{Tr} \left(\Pi_s'^2 \right) = \Pi_{\dot{\alpha}}^{\alpha} \Pi_{\alpha}^{\dot{\alpha}} = 0, \quad (2.15)$$

since,

$$\Pi_{\dot{\alpha}}^{\alpha} \Pi_{\alpha}^{\dot{\alpha}} = \epsilon_{\alpha\beta} \Pi_{\dot{\alpha}}^{\alpha} \Pi_{\dot{\beta}}^{\beta} \epsilon^{\dot{\alpha}\dot{\beta}} = \begin{cases} \lambda^2 \tilde{\lambda}_s^2, & s = 0, \pm 2, \pm 4, \dots \\ \lambda^2 \tilde{\zeta}_s^2, & s = \pm 1, \pm 3, \pm 5, \dots \end{cases}, \quad (2.16)$$

exactly as in eq(2.6). Nilpotency of $\Pi_{\dot{\alpha}}^{\alpha} \Pi_{\alpha}^{\dot{\alpha}}$, the invariant square of $(\Pi'_s)_{\dot{\alpha}}^{\alpha}$'s is then mainly due to

$$\lambda^2 = 0, \quad (2.17)$$

since for the case of anticommuting $\tilde{\zeta}_s$'s, the squares

$$\left(\tilde{\zeta}_s \right)^2 = \epsilon^{\dot{\alpha}\dot{\beta}} \tilde{\zeta}_{s\dot{\alpha}} \tilde{\zeta}_{s\dot{\beta}} = \frac{1}{2} \epsilon^{\dot{\alpha}\dot{\beta}} \left(\tilde{\zeta}_{s\dot{\alpha}} \tilde{\zeta}_{s\dot{\beta}} - \tilde{\zeta}_{s\dot{\beta}} \tilde{\zeta}_{s\dot{\alpha}} \right) \neq 0, \quad (2.18)$$

are non zero. Note that for the world line bosonic subset, we also have

$$\left(\tilde{\lambda}_s \right)^2 = 0. \quad (2.19)$$

and, in addition to $p_{\dot{\alpha}}^{\alpha} = \lambda^{\alpha} \tilde{\lambda}_{0\dot{\alpha}}$, we shall also use the following typical world line fields for our QFT model building and its dual twistor like,

$$\Pi_{\pm\dot{\alpha}}^{\alpha} = \lambda^{\alpha} \tilde{\zeta}_{\pm\dot{\alpha}}, \quad \mathcal{P}_{\pm\pm\dot{\alpha}}^{\alpha} = \lambda^{\alpha} \tilde{\lambda}_{\pm\pm\dot{\alpha}}. \quad (2.20)$$

Here $\tilde{\zeta}_{\pm\dot{\alpha}}$ and $\tilde{\lambda}_{\pm\pm\dot{\alpha}}$ are respectively anticommuting and commuting world line fermions and bosons. Note moreover the following useful properties which can help to get an overview on a possible stringy extension:

(a) In general, the above result is not only valid for quantum mechanics. These world line fields can be naturally promoted to world sheet field variables where irreducible representations of the $SO(2, R)$ rotation group of the real surface (plane) are one dimensional. The charges $\pm q$ carried by the fields are nothing but the conformal weights of 2D conformal invariance; see next subsection for technical details.

(b) Along with the set $(\Pi'_s)_{\dot{\alpha}}^{\alpha}$ eqs(2.13-2.14), we have a second family $(\Pi''_s)_{\dot{\alpha}}^{\alpha}$ where the world line spin is carried by the undotted $SO(d, 4-d)$ spinors of target space. This family is given by,

$$(\Pi''_s)_{\dot{\alpha}}^{\alpha} = (\lambda_s)^{\alpha} \tilde{\lambda}_{\dot{\alpha}}, \quad s = 0, \pm 2, \pm 4, \dots, \quad (2.21)$$

for world line bosons and by

$$(\Pi''_s)_{\dot{\alpha}}^{\alpha} = (\zeta_s)^{\alpha} \tilde{\lambda}_{\dot{\alpha}}, \quad s = \pm 1, \pm 3, \pm 5, \dots, \quad (2.22)$$

for world line fermions. The vanishing condition $\text{Tr}(\Pi''_s{}^2) = 0$ is mainly ensured by the property

$$\tilde{\lambda}^2 = \tilde{\lambda}_{\dot{\alpha}} \tilde{\lambda}^{\dot{\alpha}} = 0 \quad (2.23)$$

for all values of s .

(c) Both of the families $\Pi'_{s\dot{\alpha}}^{\alpha}$ and $\Pi''_{s\dot{\alpha}}^{\alpha}$ are in fact two special subsets of the following general one,

$$(\Pi_s)_{\dot{\alpha}}^{\alpha} = (f_{s_1})^{\alpha} \tilde{g}_{s_2\dot{\alpha}}, \quad s = s_1 + s_2. \quad (2.24)$$

Here the condition $\text{Tr}(\Pi_s^2) = 0$ is not automatically fulfilled; it is solved whenever at least one of the two world line (sheet) fields $f_{s_1}^{\alpha}$ and $\tilde{g}_{s_2\dot{\alpha}}$ is a boson.

2.2 $\Pi_{s\dot{\alpha}}^{\alpha}(\tau)$ as boundary fields

The above world line field variables, which we denote generically as $\phi = \phi(\tau)$, may be given various geometric interpretations. For instance, they may be thought of as the boundary values,

$$\phi(\tau) = \phi(\sigma^-, \sigma^+) |_{\partial\mathcal{M}} \quad (2.25)$$

of world sheet fields $\phi(\sigma^-, \sigma^+)$ of a $2d$ quantum (conformal) field theory (QFT₂). Here $\sigma^{\pm} = \frac{1}{2}(\sigma_0 \pm \sigma_1)$ are the variables parameterizing the Riemann surface $\mathcal{M}$ with boundary $\partial\mathcal{M}$ parameterized by τ . In this view, the previous $\Pi_{s\dot{\alpha}}^{\alpha}(\tau)$'s are seen as the boundary values associated with the world sheet fields,

$$\Pi_{s\dot{\alpha}}^{\alpha} = \Pi_{s\dot{\alpha}}^{\alpha}(\sigma^-, \sigma^+), \quad \Pi_{s\dot{\alpha}}^{\alpha}(\tau) = \Pi_{s\dot{\alpha}}^{\alpha}(\sigma^-, \sigma^+) |_{\partial\mathcal{M}}, \quad (2.26)$$

and s gets interpreted as the conformal spin, i.e

$$\left(\sigma^+ \frac{\partial}{\partial \sigma^+} - \sigma^- \frac{\partial}{\partial \sigma^-}\right) \Pi_{s\dot{\alpha}}^\alpha(\sigma) \sim s \Pi_{s\dot{\alpha}}^\alpha(\sigma). \quad (2.27)$$

In section 3, we will give other examples by using reductions of world sheet $n = (1, 1)$ superalgebra¹ with supersymmetric covariant derivatives D_+ and D_- ,

$$D_\mp^2 \sim \frac{\partial}{\partial \sigma^\pm} \quad (2.28)$$

One of these examples is given by interpreting the world line variable τ as the light cone coordinate,

$$\sigma^+ = \frac{1}{2}(\sigma_0 + \sigma_1), \quad (2.29)$$

parameterizing a left moving closed string or equivalently like $\sigma^- = \frac{1}{2}(\sigma_0 - \sigma_1)$ for right moving. An other remarkable example, which we develop in present study, is given by the association of the τ variable with an extra geometric dimension realizing the central charge

$$\mathcal{Z} \sim D_+ D_- + D_- D_+ \quad (2.30)$$

of the $n = (1, 1)$ superalgebra; see also eqs(3.1) to fix the ideas. More precisely we have,

$$\mathcal{Z}\phi(\tau) = \frac{\partial \phi(\tau)}{\partial \tau}. \quad (2.31)$$

Before going into technical details, let us say few words about these tools and where they will be used. This brings us to our next result.

2.3 A pure fermionic model

Part of this paper will be dealing with the derivation of a new pure fermionic twistor like model. Instead of the massless particle described by the world line field $x_\alpha^\alpha(\tau)$ and its conjugate momentum $p_\alpha^\alpha(\tau)$ with $\text{Tr}(p^2) = 0$, we consider rather the world line fermions,

$$\Upsilon_{+\dot{\alpha}}^\alpha = \Upsilon_{+\dot{\alpha}}^\alpha(\tau), \quad \tilde{\Upsilon}_{+\dot{\alpha}}^\alpha = \tilde{\Upsilon}_{+\dot{\alpha}}^\alpha(\tau), \quad (2.32)$$

together with their conjugate momenta

$$\pi_{-\dot{\alpha}}^\alpha = \pi_{-\dot{\alpha}}^\alpha(\tau), \quad \tilde{\pi}_{-\dot{\alpha}}^\alpha = \tilde{\pi}_{-\dot{\alpha}}^\alpha(\tau). \quad (2.33)$$

Obviously this can be viewed as the simplest case one may consider. Nevertheless, extensions involving much more fields is straightforward and it turns out that they lead

¹As a convention of notations, we use the letter n to refer to number of supercharges of world line (sheet) supersymmetry. Capital letter $\mathcal{N}$ is deserved for target space supersymmetry.

to extended target space supersymmetry. We shall not study this issue in this paper; comments will given in the conclusion and discussion section.

We also have the twistor like splitting,

$$\begin{aligned}\Upsilon_{+\dot{\alpha}}^{\alpha} &= \lambda^{\alpha} \tilde{\zeta}_{+\dot{\alpha}}, & \pi_{-\alpha}^{\dot{\alpha}} &= \lambda_{\alpha} \tilde{\zeta}_{-}^{\dot{\alpha}}, \\ \tilde{\Upsilon}_{+\dot{\alpha}}^{\alpha} &= \zeta_{+}^{\alpha} \tilde{\lambda}_{\dot{\alpha}}, & \tilde{\pi}_{-\alpha}^{\dot{\alpha}} &= \zeta_{-}^{\alpha} \tilde{\lambda}_{\dot{\alpha}}.\end{aligned}\quad (2.34)$$

where λ^{α} and $\tilde{\lambda}_{\dot{\alpha}}$ are the usual Penrose variable and $\zeta_{\pm}^{\alpha}$ and $\tilde{\zeta}_{\pm\dot{\alpha}}$ are anticommuting world line fermions,

$$\begin{aligned}\tilde{\zeta}_{+\dot{\alpha}} \tilde{\zeta}_{+\dot{\beta}} + \tilde{\zeta}_{+\dot{\beta}} \tilde{\zeta}_{+\dot{\alpha}} &= 0, \\ \tilde{\zeta}_{-\dot{\alpha}} \tilde{\zeta}_{+\dot{\beta}} + \tilde{\zeta}_{+\dot{\beta}} \tilde{\zeta}_{-\dot{\alpha}} &= 0, \\ \tilde{\zeta}_{-\dot{\alpha}} \tilde{\zeta}_{-\dot{\beta}} + \tilde{\zeta}_{-\dot{\beta}} \tilde{\zeta}_{-\dot{\alpha}} &= 0,\end{aligned}\quad (2.35)$$

and similar relations for undotted world line spinors. As it is known, the conjugate momenta $\pi_{\mp\alpha}^{\dot{\alpha}}$ for fermions can be also imagined as just $\epsilon_{\alpha\beta}\epsilon^{\dot{\alpha}\dot{\beta}}\Upsilon_{\mp\dot{\alpha}}^{\beta}$. By substituting the expressions of $\Upsilon_{+\dot{\alpha}}^{\alpha}$ and $\pi_{-\alpha}^{\dot{\alpha}}$ back into eqs(2.35), one finds that they satisfy as well the same anticommutation relations as the ζ 's,

$$\begin{aligned}\Upsilon_{+\dot{\alpha}}^{\alpha} \Upsilon_{+\dot{\beta}}^{\beta} + \Upsilon_{+\dot{\beta}}^{\beta} \Upsilon_{+\dot{\alpha}}^{\alpha} &= 0, \\ \pi_{-\alpha}^{\dot{\alpha}} \Upsilon_{+\dot{\beta}}^{\beta} + \Upsilon_{+\dot{\beta}}^{\beta} \pi_{-\alpha}^{\dot{\alpha}} &= 0, \\ \pi_{-\alpha}^{\dot{\alpha}} \pi_{-\beta}^{\dot{\beta}} + \pi_{-\beta}^{\dot{\beta}} \pi_{-\alpha}^{\dot{\alpha}} &= 0.\end{aligned}\quad (2.36)$$

Similar relations are also valid for $\tilde{\Upsilon}$ and $\tilde{\Pi}$. By taking the trace of these relations and their tildes, we obtain the following

$$\begin{aligned}\text{Tr}(\Upsilon_{+}^2) &= 0, & \text{Tr}(\tilde{\Upsilon}_{+}^2) &= 0 \\ \text{Tr}(\pi_{-}^2) &= 0, & \text{Tr}(\tilde{\pi}_{-}^2) &= 0 \\ \text{Tr}(\pi_{-}\Upsilon_{+}) &= 0, & \text{Tr}(\tilde{\pi}_{-}\tilde{\Upsilon}_{+}) &= 0.\end{aligned}\quad (2.37)$$

they are nothing but the field property $\text{Tr}(\Upsilon_{\pm}^2) = 0$ and partners derived in sub-section 2.1. The pure fermionic world line field action describing the dynamics of these fields is given,

$$\begin{aligned}\mathcal{S}_F[\Upsilon, \pi, \tilde{\Upsilon}, \tilde{\pi}] &= \int_{\mathbb{R}} d\tau \text{Tr} \left(\pi_{-\dot{\alpha}}^{\alpha} \frac{\partial}{\partial \tau} \Upsilon_{+\alpha}^{\dot{\alpha}} + \tilde{\pi}_{-\dot{\alpha}}^{\alpha} \frac{\partial}{\partial \tau} \tilde{\Upsilon}_{+\alpha}^{\dot{\alpha}} \right) \\ &\quad - \int_{\mathbb{R}} d\tau \text{Tr} [\eta_{--} (\Upsilon_{+}^2) + \eta_{++} (\pi_{-}^2) + \eta_{+-} (\pi_{-}\Upsilon_{+})] \\ &\quad - \int_{\mathbb{R}} d\tau \text{Tr} [\tilde{\eta}_{--} (\tilde{\Upsilon}_{+}^2) + \tilde{\eta}_{++} (\tilde{\pi}_{-}^2) + \tilde{\eta}_{+-} (\tilde{\pi}_{-}\tilde{\Upsilon}_{+})].\end{aligned}\quad (2.38)$$

where $\eta_{\pm\pm}$, η_{+-} and $\tilde{\eta}_{\pm\pm}$, $\tilde{\eta}_{+-}$ are auxiliary world line gauge fields capturing the three constraint eqs(2.37). We will see later; in particular for the case of real world line fermions, that these field constraints are captured by a $SP(2, R) \times SP(2, R)$ gauge symmetry.

2.4 World line (sheet) & target space supersymmetries

In above field action, one notes that the solution (2.34) of the constraint eqs(2.37), which can be defined also as,

$$\frac{\delta \mathcal{S}_F}{\delta \eta_{rs}} = 0, \quad \frac{\delta \mathcal{S}_F}{\delta \tilde{\eta}_{rs}} = 0, \quad r, s = \pm \quad (2.39)$$

involves two kinds of world line field pairs: (i) the world line bosons

$$\lambda^\alpha, \quad \tilde{\lambda}_{\dot{\alpha}}, \quad (2.40)$$

and (ii) the fermionic partners

$$\zeta_+^\alpha, \quad \tilde{\zeta}_{+\dot{\alpha}}. \quad (2.41)$$

If supplementing these twistor like variables by their conjugate momenta namely

$$\mu^\alpha, \quad \tilde{\mu}_{\dot{\alpha}}, \quad \zeta_-^\alpha, \quad \tilde{\zeta}_{-\dot{\alpha}}, \quad (2.42)$$

one sees that these fields altogether form world line supersymmetric multiplets as shown below,

$$(\lambda^\alpha, \zeta_+^\alpha), \quad (\tilde{\lambda}_{\dot{\alpha}}, \tilde{\zeta}_{+\dot{\alpha}}), \quad (\mu^\alpha, \zeta_-^\alpha), \quad (\tilde{\mu}_{\dot{\alpha}}, \tilde{\zeta}_{-\dot{\alpha}}). \quad (2.43)$$

This property shows that the field action (2.38) can be made supersymmetric. In this subsection, we first give preliminary results on world line (sheet) extensions of our model. Then we give the link between the pure fermionic model and target space supersymmetry.

2.4.1 World line supersymmetry

In standard Penrose construction, one uses the world line twistor field variables,

$$\lambda^\alpha = \lambda^\alpha(\tau), \quad \mu^{\dot{\alpha}} = \mu^{\dot{\alpha}}(\tau), \quad (2.44)$$

together with their companions with opposite $SO(d, 4-d)$ chiralities, $d = 2, 3, 4$,

$$\tilde{\mu}^\alpha = \tilde{\mu}^\alpha(\tau), \quad \tilde{\lambda}_{\dot{\alpha}} = \tilde{\lambda}_{\dot{\alpha}}(\tau). \quad (2.45)$$

As the intrinsic properties of these spinors depends on the value of d , the structure of the underlying QFTs is sensitive to these features. For the target space $\mathbb{R}^{(2,2)}$ for instance, the homogeneity group $SO(2, 2, R)$ is locally isomorphic $SL(2, R) \times \widetilde{SL}(2, R)$ and the above component field spinors are basically real; a specific property making the $\mathbb{R}^{(2,2)}$ analysis a little bit subtle. However, for the spaces $\mathbb{R}^{(1,3)}$ and $\mathbb{R}^4$, the corresponding rotation groups $SO(1, 3, R)$ and $SO(4, R)$ are respectively isomorphic to $SL(2, C)$ and $SU(2, C) \times SU(2, C)$ and the fundamental spinors are complex. For the example of the

$SL(2, C)$ case, the sector with $\tilde{\mu}^\alpha$ and $\tilde{\lambda}_{\dot{\alpha}}$ eq(2.45) is just the complex conjugate of $\mu^{\dot{\alpha}}$ and λ^α , i.e,

$$\overline{\mu}^\alpha = \overline{(\mu_{\dot{\alpha}})}, \quad \overline{\lambda}_\alpha = \overline{(\lambda^\alpha)}. \quad (2.46)$$

This complex nature of the twistor variables plays a crucial role in the algebraic geometry analysis of twistor space where holomorphic and antiholomorphic sectors factorise.

In general, the field $\mu^{\dot{\alpha}} = \frac{\delta S_T}{\delta \partial_\tau \tilde{\lambda}_{\dot{\alpha}}}$ (resp. $\tilde{\mu}_\alpha = \frac{\delta S_T}{\delta \partial_\tau \lambda^\alpha}$) is the conjugate momentum of λ^α (resp. $\tilde{\lambda}_{\dot{\alpha}}$),

$$\left\{ \tilde{\lambda}_{\dot{\alpha}}, \mu^{\dot{\beta}} \right\}_{TPB} = \delta_{\dot{\alpha}}^{\dot{\beta}}, \quad (2.47)$$

where $\{f, g\}_{TPB}$ stands for Poisson bracket in twistor space. To make contact with $\mathbb{R}^{(d, 4-d)}$ geometry, λ^α and $\mu^{\dot{\alpha}}$ should be compared with the usual phase space variables,

$$x^m \sim x_\alpha^{\dot{\alpha}}(\tau), \quad p_m \sim p_\alpha^\alpha(\tau) \quad (2.48)$$

with $\{x^m, p_n\}_{MPB} = \delta_n^m$ being the Poisson bracket.

In the twistor like fermions we are considering here, we have,

$$\tilde{\zeta}_{+\dot{\alpha}} = \tilde{\zeta}_{+\dot{\alpha}}(\tau), \quad \xi_-^{\dot{\alpha}} = \xi_-^{\dot{\alpha}}(\tau), \quad (2.49)$$

with $\xi_-^{\dot{\alpha}} = \frac{\delta S_{FT}}{\delta (\partial_\tau \tilde{\zeta}_{+\dot{\alpha}})}$ and where S_{FT} is the field action of the twistor like variables. We also have

$$\left\{ \tilde{\zeta}_{+\dot{\alpha}}, \xi_-^{\dot{\beta}} \right\}_{TGFB} = \delta_{\dot{\alpha}}^{\dot{\beta}} \quad (2.50)$$

where now $\{f_+, g_-\}_{TGFB}$ is the graded Poisson bracket in twistor space. These fermionic variables $\tilde{\zeta}_{+\dot{\alpha}}$ and $\xi_-^{\dot{\alpha}}$ should be thought of as the twistor dual to the world line fermions in 4- vector of $SO(d, 4-d)$,

$$\begin{aligned} \Upsilon_+^m &\sim \Upsilon_{+\alpha}^{\dot{\alpha}}(\tau), & \pi_{-m} &\sim \pi_{-\dot{\alpha}}^\alpha(\tau) \\ \tilde{\Upsilon}_+^m &\sim \tilde{\Upsilon}_{+\alpha}^{\dot{\alpha}}(\tau), & \tilde{\pi}_{-m} &\sim \tilde{\pi}_{-\dot{\alpha}}^\alpha(\tau) \end{aligned} \quad (2.51)$$

If forgetting for a while about the world line variable τ , one remarks that the combination of eqs(2.48-2.51) describe pairs $(x, \Upsilon_+, \tilde{\Upsilon}_+)$ and $(p, \pi_-, \tilde{\pi}_-)$, which can be viewed as world line supermultiplets,

$$\left(x^m, \Upsilon_+^m, \tilde{\Upsilon}_+^m \right), \quad (p_m, \pi_{-m}, \tilde{\pi}_{-m}), \quad m = 0, 1, 2, 3. \quad (2.52)$$

A similar conclusion may be done regarding the supermultiplets built out of twistor like variables $\lambda^\alpha, \tilde{\zeta}_{+\dot{\alpha}}, \mu^{\dot{\beta}}, \xi_-^{\dot{\beta}}$ and their partners involving the spinors with opposite chirality.

Using the representations,

$$\begin{aligned} p_m &\sim p_\alpha^\alpha = \lambda^\alpha \tilde{\lambda}_{\dot{\alpha}}, \\ \pi_{-m} &\sim \pi_{-\dot{\alpha}}^\alpha = \zeta_-^\alpha \tilde{\lambda}_{\dot{\alpha}}, \\ \tilde{\pi}_{-m} &\sim \tilde{\pi}_{-\dot{\alpha}}^\alpha = \lambda^\alpha \tilde{\zeta}_{-\dot{\alpha}}, \end{aligned} \quad (2.53)$$

it is not difficult to check that world line supersymmetric transformations,

$$\begin{aligned}\delta p_{\dot{\alpha}}^{\alpha} &\sim \varepsilon_+ \pi_{-\dot{\alpha}}^{\alpha} + \tilde{\varepsilon}_+ \tilde{\pi}_{-\dot{\alpha}}^{\alpha}, \\ \delta \pi_{-\dot{\alpha}}^{\alpha} &\sim \varepsilon_+ \frac{dp_{\dot{\alpha}}^{\alpha}}{d\tau}, \quad \delta \tilde{\pi}_{-\dot{\alpha}}^{\alpha} \sim \tilde{\varepsilon}_+ \frac{d\tilde{p}_{\dot{\alpha}}^{\alpha}}{d\tau}\end{aligned}\tag{2.54}$$

translate as well to the twistor like space. By substitution, we get $\delta p_{\dot{\alpha}}^{\alpha} = (\delta \lambda^{\alpha}) \tilde{\lambda}_{\dot{\alpha}} + \lambda^{\alpha} (\delta \tilde{\lambda}_{\dot{\alpha}})$ which should be identified with $(\varepsilon_+ \zeta_{-}^{\alpha}) \tilde{\lambda}_{\dot{\alpha}} + \lambda^{\alpha} (\tilde{\varepsilon}_+ \tilde{\zeta}_{-\dot{\alpha}})$. Thus we have,

$$\begin{aligned}\delta \lambda^{\alpha} &\sim \varepsilon_+ \zeta_{-}^{\alpha}, \quad \delta \tilde{\lambda}_{\dot{\alpha}} \sim \tilde{\varepsilon}_+ \tilde{\zeta}_{-\dot{\alpha}}, \\ \delta \zeta_{-}^{\alpha} &\sim \varepsilon_+ \frac{d\lambda^{\alpha}}{d\tau}, \quad \delta \tilde{\zeta}_{-\dot{\alpha}} \sim \tilde{\varepsilon}_+ \frac{d\tilde{\lambda}_{\dot{\alpha}}}{d\tau}.\end{aligned}\tag{2.55}$$

The pure fermionic construction to be developed in this paper may be then viewed as a sub-sector of the odd part of a super-world line extension of Penrose model.

2.4.2 World sheet supersymmetry

Following the view given by eq(2.25), the above super-world line field model can be embedded in turn in world sheet $n = (1, 1)$ supersymmetric field theory with target space $\mathbb{R}^{(d, 4-d)}$. There, generic fields ϕ are functions of the world sheet variables $\sigma = (\sigma^-, \sigma^+)$. The field variables,

$$\begin{aligned}x_{\dot{\alpha}}^{\alpha} &= x_{\dot{\alpha}}^{\alpha}(\sigma), \quad \Upsilon_{\pm\dot{\alpha}}^{\alpha} = \Upsilon_{\pm\dot{\alpha}}^{\alpha}(\sigma), \\ p_{\dot{\alpha}}^{\alpha} &= p_{\dot{\alpha}}^{\alpha}(\sigma), \quad \pi_{\pm\dot{\alpha}}^{\alpha} = \pi_{\pm\dot{\alpha}}^{\alpha}(\sigma),\end{aligned}\tag{2.56}$$

and the corresponding twistor like variables,

$$\begin{aligned}\lambda^{\alpha} &= \lambda^{\alpha}(\sigma), \quad \zeta_{\pm}^{\alpha} = \zeta_{\pm}^{\alpha}(\sigma), \\ \mu^{\dot{\alpha}} &= \mu^{\dot{\alpha}}(\sigma), \quad \xi_{\mp}^{\dot{\alpha}} = \xi_{\mp}^{\dot{\alpha}}(\sigma), \\ \tilde{\lambda}_{\dot{\alpha}} &= \tilde{\lambda}_{\dot{\alpha}}(\sigma), \quad \tilde{\zeta}_{\pm\dot{\alpha}} = \tilde{\zeta}_{\pm\dot{\alpha}}(\sigma) \\ \tilde{\mu}_{\alpha} &= \tilde{\mu}_{\alpha}(\sigma), \quad \tilde{\xi}_{\mp\alpha} = \tilde{\xi}_{\mp\alpha}(\sigma),\end{aligned}\tag{2.57}$$

form world sheet supermultiplets. We also have the following graded Poisson brackets,

$$\begin{aligned}\{x_{\dot{\alpha}}^{\alpha}(\sigma), p_{\dot{\beta}}^{\beta}(\sigma')\} &= \delta_{\dot{\beta}}^{\alpha} \delta_{\dot{\alpha}}^{\beta} \delta(\sigma - \sigma') \\ \{\lambda^{\alpha}(\sigma), \tilde{\mu}_{\dot{\beta}}(\sigma')\} &= \delta_{\dot{\beta}}^{\alpha} \delta(\sigma - \sigma') \\ \{\tilde{\lambda}_{\dot{\alpha}}(\sigma), \mu^{\dot{\beta}}(\sigma')\} &= \delta_{\dot{\alpha}}^{\dot{\beta}} \delta(\sigma - \sigma') \\ \{\zeta_{\pm}^{\alpha}(\sigma), \tilde{\xi}_{\mp\dot{\beta}}(\sigma')\} &= \delta_{\dot{\beta}}^{\alpha} \delta(\sigma - \sigma') \\ \{\tilde{\zeta}_{\pm\dot{\alpha}}(\sigma), \xi_{\mp}^{\dot{\beta}}(\sigma')\} &= \delta_{\dot{\alpha}}^{\dot{\beta}} \delta(\sigma - \sigma'),\end{aligned}\tag{2.58}$$

and all remaining others are zero. Typical world sheet supersymmetric transformations are given by,

$$\begin{aligned}\delta_\varepsilon x_\alpha^\alpha(\sigma) &\sim i\varepsilon_+ \Upsilon_{-\dot{\alpha}}^\alpha(\sigma) + i\varepsilon_- \Upsilon_{+\dot{\alpha}}^\alpha(\sigma) \\ \delta_\varepsilon \Upsilon_{\pm\dot{\alpha}}^\alpha(\sigma) &\sim \varepsilon_\mp \partial_{\pm\pm} x_\alpha^\alpha(\sigma).\end{aligned}\quad (2.59)$$

$\varepsilon_\mp$ are the parameters of the $n = (1, 1)$ superalgebra to be studied in next section. Similar transformations may be written down for $(p_\alpha^\alpha(\sigma), \pi_{-\dot{\alpha}}^\alpha(\sigma))$ and the corresponding twistor like ones.

2.4.3 Target space supersymmetry

The result we give here is that the quantum spectrum of the pure fermionic with field action $\mathcal{S}_F[\Upsilon, \pi, \tilde{\Upsilon}, \tilde{\pi}]$ eqs(2.38) has $4D$ target space supersymmetry. To derive $4D$ $\mathcal{N} = 1$ target space supersymmetry, consider the following particular twistor like fermionic world line field action,

$$\begin{aligned}\mathcal{S}_{FT} &\sim \int d\tau \left[\tilde{\mu}_\alpha \partial_\tau \lambda^\alpha - \xi_{-\dot{\alpha}}^\alpha \partial_\tau \tilde{\zeta}_{+\dot{\alpha}} \right] \\ &\quad - \int d\tau \left[\mu^{\dot{\alpha}} \partial_\tau \tilde{\lambda}_{\dot{\alpha}} - \tilde{\xi}_{-\alpha} \partial_\tau \zeta_{+\alpha} \right].\end{aligned}\quad (2.60)$$

This action has a set of symmetries to be studied in section 5 and lead to $\mathcal{N} = 1$ supersymmetric quantum spectrum in target space coordinated by $\{\lambda^\alpha, \mu^{\dot{\alpha}}\}$. Eq(2.60) involves two world line fermions ζ_-^α and $\mu^{\dot{\alpha}}$ together with their conjugate momenta; more world line spinors leads to $4D$ target space extended supersymmetry.

Target space supersymmetry is obtained by quantizing the system. This is achieved by promoting the graded Poisson brackets the phase space variables,

$$\begin{aligned}\left\{ \tilde{\lambda}_{\dot{\alpha}}, \mu^{\dot{\beta}} \right\}_{GPB} &\sim \delta_{\dot{\alpha}}^{\dot{\beta}}, & \left\{ \lambda^\alpha, \tilde{\mu}_\beta \right\}_{GPB} &\sim \delta_\beta^\alpha, \\ \left\{ \tilde{\zeta}_{+\dot{\alpha}}, \xi_{-\dot{\beta}} \right\}_{GPB} &\sim \delta_{\dot{\alpha}}^{\dot{\beta}}, & \left\{ \zeta_{+\alpha}^\alpha, \tilde{\xi}_{-\beta} \right\}_{GPB} &\sim \delta_\beta^\alpha,\end{aligned}\quad (2.61)$$

into graded canonical commutation relations.

$$\begin{aligned}\left[\tilde{\lambda}_{\dot{\alpha}}, \mu^{\dot{\beta}} \right]_- &\sim \delta_{\dot{\alpha}}^{\dot{\beta}}, & \left[\lambda^\alpha, \tilde{\mu}_\beta \right]_- &\sim \delta_\beta^\alpha, \\ \left[\tilde{\zeta}_{+\dot{\alpha}}, \xi_{-\dot{\beta}} \right]_+ &\sim \delta_{\dot{\alpha}}^{\dot{\beta}}, & \left[\zeta_{+\alpha}^\alpha, \tilde{\xi}_{-\beta} \right]_+ &\sim \delta_\beta^\alpha.\end{aligned}\quad (2.62)$$

Here $[,]_-$ and $[,]_+$ (to avoid confusion with graded Poisson bracket $\{ , \}_{GPB}$) stand for commutator and anticommutator respectively. Then solve these relations like

$$\begin{aligned}\mu^{\dot{\alpha}} &\sim \frac{\partial}{\partial \tilde{\lambda}_{\dot{\alpha}}}, & \tilde{\mu}_\alpha &\sim \frac{\partial}{\partial \lambda^\alpha}, \\ \xi_{-\dot{\alpha}}^\alpha &\sim \frac{\partial}{\partial \tilde{\zeta}_{+\dot{\alpha}}}, & \tilde{\xi}_{+\alpha} &\sim \frac{\partial}{\partial \zeta_{-\alpha}^\alpha}.\end{aligned}\quad (2.63)$$

In this representation, the conjugate momenta are then given by differential operators which act on the wave functions Φ living on a graded space as shown below,

$$\Phi = \Phi(\lambda^\alpha, \mu^{\dot{\alpha}} | \zeta_+^\alpha, \xi_-^{\dot{\alpha}}). \quad (2.64)$$

Since ζ_+^α and $\xi_-^{\dot{\alpha}}$ are fermions, the waves Φ can be thought of as superfields on the superspace $\mathbb{R}^{(d,4-d|4)}$; and moreover can be expanded as usual into a finite series of power of ζ_-^α and $\xi_-^{\dot{\alpha}}$. These superfields define just off shell representations of the four dimensional $\mathcal{N} = 1$ supersymmetry,

$$\begin{aligned} \{D_{+\alpha}, D_-^{\dot{\alpha}}\} &= 2 P_\alpha^{\dot{\alpha}}, \\ [P_\alpha^{\dot{\alpha}}, D_{+\alpha}] &= [P_\alpha^{\dot{\alpha}}, D_-^{\dot{\alpha}}] = 0. \end{aligned} \quad (2.65)$$

In twistor like variables, this superalgebra can be realized as,

$$\begin{aligned} D_{+\dot{\alpha}} &= \frac{\partial}{\partial \xi_-^{\dot{\alpha}}} + \zeta_+^\alpha \lambda_\alpha \frac{\partial}{\partial \mu^{\dot{\alpha}}}, \\ D_{-\alpha} &= \frac{\partial}{\partial \zeta_+^\alpha} + \xi_-^{\dot{\alpha}} \mu_{\dot{\alpha}} \frac{\partial}{\partial \lambda^\alpha}, \\ P_\alpha^{\dot{\alpha}} &= \lambda_\alpha \frac{\partial}{\partial \mu_{\dot{\alpha}}} + \mu^{\dot{\alpha}} \frac{\partial}{\partial \lambda^\alpha}. \end{aligned} \quad (2.66)$$

To fix the ideas, let us focus on $\mathbb{R}^{(1,3|4)}$ superspace and give the example of $4D$ $\mathcal{N} = 1$ chiral multiplet; similar analysis can be done for gauge multiplet. In this case, $\tilde{\lambda}_\alpha = \bar{\lambda}_\alpha$, $\tilde{\zeta}_+^{\dot{\alpha}} = \bar{\zeta}_+^{\dot{\alpha}}$ and eq(2.65) has a chiral representation. The superfield Φ has no dependence in $\bar{\xi}_-^{\dot{\alpha}}$ and its expansion reads as follows,

$$\Psi(\lambda, \mu | \zeta_+^\alpha) = \phi(\lambda, \mu) + \zeta_+^\alpha \psi_{-\alpha}(\lambda, \mu) + \zeta_+^2 F_{--}(\lambda, \mu). \quad (2.67)$$

The component fields $(\phi, \psi_\alpha, F_{--})$ are nothing but the scalar multiplet of four dimensional $\mathcal{N} = 1$ superalgebra(2.65). For other supersymmetric representations and explicit details for the target spaces $\mathbb{R}^{(d,4-d)}$, $d = 2, 3, 4$, see section 5 and eqs(5.57,5.58,5.59).

3 Embedding world line models in 2D superalgebra

Because of geometric interpretation purpose and for later use, we start this section by discussing briefly the general superspace realization of the world sheet $\mathfrak{n} = (1, 1)$ supersymmetric algebra. Then we consider particular cases associated with some special subalgebras of this $\mathbb{Z}_2$ graded algebra. This helps us to derive the symmetry properties of the world line fields of the twistor like fermion model.

Roughly, the $\mathfrak{n} = (1, 1)$ superalgebra lives on Riemann surfaces $\mathcal{M}$ (string world sheet) and can, roughly, be defined on a local chart of $\mathcal{M}$ as,

$$\begin{aligned}
\{D_+, D_+\} &= -2a\partial_{++}, \\
\{D_-, D_-\} &= -2b\partial_{--} \\
\{D_\pm, D_\mp\} &= -2c\mathcal{Z} \\
[\partial_{\pm\pm}, Q_\pm] &= [\partial_{\pm\pm}, \mathcal{Z}] = [\mathcal{Z}, Q_\pm] = 0,
\end{aligned} \tag{3.1}$$

In these relations, the $D_\pm$'s are the usual supersymmetric covariant derivatives of the superalgebra and $\mathcal{Z}$ is its unique central charge. The latter is ignored in almost all simple applications; but in a complete analysis it should be taken into account as plays a crucial role in the analysis of BPS states and non perturbative dynamics. The extra parameters a, b and c are real numbers used as a trick to recover particular limits. Note in passing that for the case of $\mathfrak{n} = (1, 1)$ superalgebra, these extra parameters should be non zero and can be taken as

$$a = b = c = 1. \tag{3.2}$$

Below we want to consider as well particular cases associated with some special limits which are captured by the singular values of these parameters. Of particular interest is the set of cases encoded collectively by the solution of the relation,

$$abc = 0. \tag{3.3}$$

Representations involving these cases are related to world line supersymmetric representations and boundary conformal field theory. Let us give below some details.

3.1 Superspace realization

Using standard results on supersymmetric field theories, it is not difficult to see that the superspace representation of the above superalgebra is given by the following,

$$\begin{aligned}
D_+ &= \partial_+ - a\theta_-\partial_{++} - c\theta_+\frac{\partial}{\partial\varsigma}, \\
D_- &= \partial_- - b\theta_+\partial_{--} - c\theta_-\frac{\partial}{\partial\varsigma} \\
\partial_\pm &= \frac{\partial}{\partial\theta^\mp}, \quad \partial_{\pm\pm} = \frac{\partial}{\partial\sigma^\mp}, \quad \mathcal{Z} = \frac{\partial}{\partial\varsigma}.
\end{aligned} \tag{3.4}$$

These are superspace differential operators acting on superfields

$$\Phi = \Phi(\sigma, \varsigma, \theta) \tag{3.5}$$

living on the $(3|2)$ dimensional real superspace $\mathbb{R}^{(3|2)}$ with supercoordinates,

$$Z = (\sigma^+, \sigma^-, \varsigma \mid \theta^+, \theta^-) \tag{3.6}$$

The parameters $\sigma^+, \sigma^-, \varsigma$ are commuting variables ($\mathbb{C}$ - numbers) and $\theta^\pm \sim \theta_\mp$ are Majorana Grassmann coordinates fulfilling

$$\begin{aligned}\{\theta_\pm, \theta_\mp\} &= 0, & \{\partial_\mp, \theta_\pm\} &= 1 \\ [\sigma^\pm, \theta_\pm] &= 0, & [\varsigma, \theta_\pm] &= 0.\end{aligned}\quad (3.7)$$

Note that ς is an extra real variable realizing the charge $\mathcal{Z}$ as a translation operator in same way as do $\sigma^\mp$ for energy momentum vector $\mathcal{P}_{\pm\pm}$. The latters are realized as space time translations $\partial_{\pm\pm}$. Typical hermitian superfields $\Phi_s(\sigma, \varsigma, \theta^\pm)$ have the following θ -expansion,

$$\Phi_s(\sigma, \varsigma, \theta^\pm) = \phi_s(\sigma, \varsigma) + i\theta_- \psi_{s+1}(\sigma, \varsigma) + i\theta_+ \psi_{s-1}(\sigma, \varsigma) + i\theta_- \theta_+ D_s(\sigma, \varsigma). \quad (3.8)$$

For the special case $s = 0$, the scalar superfield Φ contains two real bosonic fields, ϕ and D and two Majorana fermions $\psi_\pm$. We also have the expansions,

$$\begin{aligned}D_+ \Phi &= i\psi_+ - a\theta_- \partial_{++}\phi + \theta_+ \left(iD - c \frac{\partial \phi}{\partial \varsigma} \right) + i\theta_- \theta_+ \left(c \frac{\partial \psi_+}{\partial \varsigma} - a \partial_{++}\psi_- \right), \\ D_- \Phi &= i\psi_- - b\theta_+ \partial_{--}\phi - \theta_- \left(iD + c \frac{\partial \phi}{\partial \varsigma} \right) - i\theta_- \theta_+ \left(c \frac{\partial \psi_-}{\partial \varsigma} - b \partial_{--}\psi_+ \right),\end{aligned}\quad (3.9)$$

which can be used to build the supersymmetric component field action, see also eq(3.18). Note that we have used,

$$(i\theta_- \theta_+)^\dagger = i\theta_- \theta_+ \quad (3.10)$$

By making appropriate choices of the parameters a, b and c in eq(3.4), one distinguishes several cases to recover world line supersymmetry. Besides the case where D_- and D_+ and $\partial_{--}, \partial_{++}, \frac{\partial}{\partial \varsigma}$ are identified to D and $\frac{d}{dt}$, i.e

$$D^2 = \frac{d}{dt}, \quad \left[D, \frac{d}{dt} \right] = 0, \quad (3.11)$$

we have moreover the three following word line like sub-superalgebras of eqs(3.1):

(i) Case $a = c = 0, b \neq 0$

This particular case corresponds to the so called $(0, 1)$ heterotic supersymmetry generated by $\{D_-, \partial_{--}\}$, i.e

$$\begin{aligned}\{D_+, D_+\} &= 0, \\ \{D_-, D_-\} &= -2b\partial_{--} \\ \{D_+, D_-\} &= 0\end{aligned}\quad (3.12)$$

and can be realized on the (right moving) world line superspace $\mathbb{R}^{(1|1)}$ parameterized by $(\sigma^- | \theta^-)$. We have

$$D_+ = \partial_+, \quad D_- = \partial_- - b\theta_+ \partial_{--}, \quad (3.13)$$

with $\theta_+^2 = 0$.

(ii) Case $b = c = 0, a \neq 0$

This situation corresponds to $(1, 0)$ heterotic supersymmetry

$$\{D_+, D_+\} = -2a\partial_{++}, \quad (3.14)$$

generated by D_+ and ∂_{++} and is realized on a (left moving) world line superspace $\mathbb{R}^{(1|1)}$ as well. The structure of the supersymmetric field theory based on this case is analogous to the previous one.

(iii) Case $a = b = 0, c \neq 0$

In this situation, there is no "light cone" energy momentum,

$$\mathcal{P}_{++} = \mathcal{P}_{--} = 0, \quad (3.15)$$

but we do have a non zero central charge contribution. The corresponding world line superalgebra

$$\begin{aligned} \{D_+, D_+\} &= 0, \\ \{D_-, D_-\} &= 0 \\ \{D_\pm, D_\mp\} &= -2c\mathcal{Z}. \end{aligned} \quad (3.16)$$

This special sub-superalgebra of (3.1) generated by $\{D_\pm, \mathcal{Z}\}$ and is realized on a world line superspace $\mathbb{R}^{(1|2)}$ parameterized by $(\varsigma \mid \theta^+, \theta^-)$ as follows,

$$\begin{aligned} D_+ &= \partial_+ - c\theta_+ \frac{\partial}{\partial \varsigma}, \\ D_- &= \partial_- - c\theta_- \frac{\partial}{\partial \varsigma} \\ \mathcal{Z} &= \frac{\partial}{\partial \varsigma}. \end{aligned} \quad (3.17)$$

Here there is no dependence in $\sigma^\pm$ variables.

3.2 Superfield action

According to which kind of the above sub-symmetries one is interested in, we can write down various world line type supersymmetric field actions. These superfield models should, a priori, be also derived by starting from the following parent superspace action

$$S \sim \int_{\mathbb{R}^{(3|2)}} d^2\sigma d\varsigma d^2\theta \left[\frac{1}{2} D_- \Phi D_+ \Phi + iW(\Phi) \right]. \quad (3.18)$$

Then make appropriate reductions as outlined above by keeping track with the implementation of the constraint eqs to reconstitue missing parts in kinetic terms of $(1, 0)$ and

(0, 1) heterotic models. In eq(3.18), $W(\Phi)$ is the superpotential; in general given by a polynom in Φ ; but here we shall consider a free field theory and so we can ignore it.

Putting the expansion (3.8) back into this superfield action, we get after integrating out the Grassmann variables $\theta_{\pm}$ the following component fields action

$$\begin{aligned}
S \sim & \int_{\mathbb{R}^3} d^2\sigma d\varsigma \left[\frac{ab}{2} \partial_{--}\phi \partial_{++}\phi + \frac{c^2}{2} \left(\frac{\partial\phi}{\partial\varsigma} \right)^2 - \frac{1}{2} D^2 \right] \\
& + \int_{\mathbb{R}^3} d^2\sigma d\varsigma \left[\frac{a}{2} \psi_- \partial_{++}\psi_- - \frac{b}{2} \psi_+ \partial_{--}\psi_+ \right] \\
& + \int_{\mathbb{R}^3} d^2\sigma d\varsigma \left[\frac{c}{2} \left(\psi_+ \frac{\partial}{\partial\varsigma} \psi_- - \psi_- \frac{\partial}{\partial\varsigma} \psi_+ \right) \right],
\end{aligned} \tag{3.19}$$

where we have exhibited the dependence in the a, b and c moduli. Note that for the particular case where

$$a = b = 0, \quad c = 1 \tag{3.20}$$

the above action reduces, roughly, to the following world line one,

$$\begin{aligned}
S_{reduced} [\phi, \psi_{\pm}] \sim & \int_{\mathbb{R}} d\varsigma \left[\frac{1}{2} \left(\frac{\partial\phi}{\partial\varsigma} \right)^2 - \frac{1}{2} D^2 \right] \\
& + \int_{\mathbb{R}} d\varsigma \left[\frac{1}{2} \left(\psi_+ \frac{\partial}{\partial\varsigma} \psi_- - \psi_- \frac{\partial}{\partial\varsigma} \psi_+ \right) \right],
\end{aligned} \tag{3.21}$$

where now $\phi = \phi(\varsigma)$ and $\psi_{\pm} = \psi_{\pm}(\varsigma)$. Note also that as expected, the auxiliary field D is non dynamical and so can be eliminated through its equation of motion, an operation once done leads to an on-shell supersymmetric field action.

Furthermore letting the supercharges act on the superfield Φ , we can immediately get the infinitesimal supersymmetric transformations,

$$\begin{aligned}
\delta\phi & \sim i\varepsilon_- \psi_+ + i\varepsilon_+ \psi_- \\
\delta\psi_+ & \sim ia\varepsilon_- \partial_{++}\phi + \varepsilon_+ \left(iD + c \frac{\partial\phi}{\partial\varsigma} \right) \\
\delta\psi_- & \sim ib\varepsilon_+ \partial_{--}\phi - \varepsilon_- \left(iD - c \frac{\partial\phi}{\partial\varsigma} \right) \\
\delta D & \sim \varepsilon_- \left(c \frac{\partial\psi_+}{\partial\varsigma} - a \partial_{++}\psi_- \right) - i\varepsilon_+ \left(c \frac{\partial\psi_-}{\partial\varsigma} - b \partial_{--}\psi_+ \right)
\end{aligned} \tag{3.22}$$

With these tools at hand, we turn now to the main objectif of the present study.

4 Fermionic like model

First we consider the pure bosonic case where the role of the generic field ϕ will be played by the target space time coordinates x^m and their conjugate momenta p_m . Then,

we study the derivation of the pure fermionic analog of the Penrose twistor model. In this case, the role of world line fermions $\psi_{\pm}$ is played by the four components world line field $\Upsilon_{\pm}^m$.

4.1 Standard twistor model: Review and comments

Roughly, twistor target space formalism allows an alternative description of the dynamics of a free massless bosonic particle with 4- energy momentum vector p_m satisfying the usual massless condition $\eta^{mn}p_m p_n = 0$ with η^{mn} the (inverse) metric of the real $\mathbb{R}^{(d,4-d)}$ space. As η_{mn} may have three possible configuration in one to one with the values

$$d = 0, 1, 2, \quad (4.1)$$

we will discuss a special example namely the case of Minkowski space $\mathbb{R}^{(1,3)}$. But the formalism we will give can be also done for the two other cases: $\mathbb{R}^4$ and $\mathbb{R}^{(2,2)}$. To derive the twistor gauge model from the usual Minkowski space formulation, we have to proceed in steps. The key idea can be summarized as follows:

4.1.1 bi-spinors

The first step consists on using $SO(1,3) \simeq SL(2,C)$ isomorphism to express 4-vectors of $\mathbb{R}^{(1,3)}$ in terms of $SL(2,C)$ spinors. Instead of phase space coordinates x^m and p_m , which can be also written as $(\frac{1}{2}, \frac{1}{2})$ bi-spinors, twistor formalism uses rather

$$\left(\frac{1}{2}, 0\right), \quad \text{and} \quad \left(0, \frac{1}{2}\right), \quad (4.2)$$

spinor fields as fundamental field variables. Before going through technical details, note first the following useful features:

(i) Given two spinors λ_1^α and λ_2^β , $\alpha, \beta = 1, 2$, of positive chirality,

$$\lambda^\alpha \sim \left(\frac{1}{2}, 0\right), \quad (4.3)$$

we define the following $SL(2,C)$ invariant

$$\lambda_1 \cdot \lambda_2 = \epsilon_{\alpha\beta} \lambda_1^\alpha \lambda_2^\beta, \quad \epsilon_{\alpha\beta} = -\epsilon_{\beta\alpha} \quad (4.4)$$

Similarly, given two spinors $\bar{\lambda}_{\dot{\alpha}1}$ and $\bar{\lambda}_{\dot{\beta}2}$ of negative chirality transforming in $(0, \frac{1}{2})$ representation, we can build a second $SL(2,C)$ invariant using dotted variables,

$$\bar{\lambda}_1 \cdot \bar{\lambda}_2 = \epsilon^{\dot{\alpha}\dot{\beta}} \bar{\lambda}_{\dot{\alpha}1} \bar{\lambda}_{\dot{\beta}2}, \quad \overline{(\epsilon^{\alpha\beta})} = \epsilon_{\dot{\beta}\dot{\alpha}}. \quad (4.5)$$

For the $SO(3,1)$ case, eqs(4.4) and (4.5) are related by usual complex conjugation ($\overline{(\lambda^\alpha)} = \bar{\lambda}_{\dot{\alpha}}$ and so on); but in the general $SO(p,q)$ case ($p+q=4$), one has to implement

moreover the special properties of the corresponding spinors.

(ii) Using $SL(2, C)$ spinors, energy momentum vector p_m may be represented line in eq(2.4). In bi-spinor notations, we have

$$p_{\dot{\beta}}^{\alpha} = \lambda^{\alpha} \bar{\lambda}_{\dot{\beta}}. \quad (4.6)$$

Note that the correspondence

$$p_m \quad \longleftrightarrow \quad (\lambda, \bar{\lambda})$$

is not a one to one; since the gauge change

$$\begin{aligned} \lambda^{\alpha} &\longrightarrow e^{i\theta} \lambda^{\alpha} \\ \bar{\lambda}_{\dot{\beta}} &\longrightarrow e^{-i\theta} \bar{\lambda}_{\dot{\beta}} \end{aligned} \quad (4.7)$$

with $\theta \in \mathbb{R}$, leaves eq(4.6) invariant. This symmetry will play a crucial role in the Penrose twistor model.

4.1.2 Twistor gauge theory

In the second step we introduce Penrose variables $(\lambda^{\alpha}, \mu^{\dot{\alpha}})$ parameterizing the projective twistor space $PT = CP^3$. Their complex conjugates $(\bar{\mu}_{\alpha}, \bar{\lambda}_{\dot{\alpha}})$ parameterize the anti-holomorphic sector. The coordinates $(\lambda^{\alpha}, \mu^{\dot{\alpha}})$ are commuting phase space variables taken in the fundamental representations of the $SL(4, C)$ twistor group. The isodoublet λ^{α} is as before while

$$\mu^{\dot{\alpha}} = \frac{\delta \mathcal{S}_{twistor}}{\delta (\partial_{\tau} \bar{\lambda}_{\dot{\alpha}})} \quad (4.8)$$

is the conjugate momentum of $\bar{\lambda}_{\dot{\alpha}}$ with $\mathcal{S}_{twistor}$ being the twistor field action to be given later on. Quantum mechanically, the variable $\mu^{\dot{\beta}}$ gets promoted to an operator $\mu^{\dot{\beta}}$ satisfying the canonical commutation relation,

$$[\bar{\lambda}_{\dot{\alpha}}, \mu^{\dot{\beta}}] = \delta_{\dot{\alpha}}^{\dot{\beta}} \quad (4.9)$$

and may be defined as,

$$\mu^{\dot{\alpha}} = -\frac{\partial}{\partial \bar{\lambda}_{\dot{\alpha}}}. \quad (4.10)$$

We will turn later to the implication of these quantum equations, for the moment note that in the classical setting, the geometric points $x_{\dot{\beta}}^{\dot{\alpha}}$ of the real 4D Minkowski space-time are associated with the so-called incidence relation,

$$\mu^{\dot{\alpha}} = i x_{\dot{\beta}}^{\dot{\alpha}} \lambda^{\dot{\beta}}, \quad \bar{\mu}_{\alpha} = i \bar{\lambda}_{\dot{\beta}} x_{\alpha}^{\dot{\beta}}, \quad \overline{(x_{\dot{\beta}}^{\dot{\alpha}})} = x_{\alpha}^{\dot{\beta}}, \quad (4.11)$$

describing a line in the spinor space with "slope" $(x_{\beta}^{\dot{\alpha}})$. These eqs may be also viewed as the solution of the twistor constraint eq,

$$\bar{\lambda}_{\dot{\alpha}}\mu^{\dot{\alpha}} - \lambda^{\alpha}\bar{\mu}_{\alpha} = 0, \quad (4.12)$$

describing a helicity zero massless particle. To see this property, it is interesting to denote collectively the twistor variables as

$$Z^a = (\lambda^{\alpha}, \mu^{\dot{\alpha}}), \quad (4.13)$$

with index $a = \alpha, \dot{\alpha}$ and projective symmetry as

$$Z^a \longrightarrow Z'^a = e^{i\theta} Z^a \quad (4.14)$$

Rising and lowering index a can be done by help of the metric

$$\Sigma_{ab} = \begin{pmatrix} 0 & \epsilon_{\dot{\alpha}\dot{\beta}} \\ \epsilon_{\alpha\beta} & 0 \end{pmatrix}, \quad \Sigma_{ab}\Sigma^{bc} = \delta_a^c. \quad (4.15)$$

For instance, we have

$$Z_a = \Sigma_{ab}Z^b = (\mu_{\dot{\alpha}}, \lambda_{\alpha}). \quad (4.16)$$

Note that the twistor variable Z^a and its complex conjugate

$$(Z^a)^{\dagger} = \bar{Z}_a = (-\bar{\mu}_{\dot{\alpha}}, \bar{\lambda}_{\alpha}) \quad (4.17)$$

are respectively in 4 and $\bar{4}$ representations of the non compact group $SU(2, 2, C) \simeq SO(4, 2, \mathbb{R})$. The latter is the conformal symmetry of the massless (spinless) relativistic particle in $4D$ space-time and appears as a hidden invariance in the underlying Minkowski space lagrangian density. Notice moreover that using eqs(4.11), the following $SU(2, 2)$ invariant

$$\bar{Z}_a Z^a = \bar{\lambda}_{\dot{\alpha}}\mu^{\dot{\alpha}} - \lambda^{\alpha}\bar{\mu}_{\alpha} \quad (4.18)$$

vanishes identically

$$\bar{Z}_a Z^a = 0 \quad (4.19)$$

This constraint eq, which will be derived rigorously later on, can be checked directly by substituting directly $\mu^{\dot{\alpha}}$ and $\bar{\mu}_{\alpha}$ by their expressions (4.11). We have $\bar{\lambda}_{\dot{\alpha}}\mu^{\dot{\alpha}} = i(\bar{\lambda}_{\dot{\alpha}}x_{\beta}^{\dot{\alpha}}\lambda^{\beta})$ which compensates exactly $\bar{\mu}_{\alpha}\lambda^{\alpha} = i(\bar{\lambda}_{\beta}x_{\alpha}^{\dot{\beta}}\lambda^{\alpha})$.

To derive eq(4.19), we have to identify first the twistor field action $\mathcal{S}_{twistor} = \mathcal{S}[Z, \bar{Z}]$. By solving $\mu^{\dot{\alpha}} = \frac{\delta \mathcal{S}_{twistor}}{\delta \tau \lambda_{\dot{\alpha}}}$ eq(4.9) and its complex conjugate and using eqs(4.13), it is not difficult to see that the manifestly $SU(2, 2)$ invariant $\mathcal{S}_{twistor}$ reads as follows,

$$\mathcal{S}_{twistor}[Z, \bar{Z}] = - \int d\tau \left[\bar{Z}_a i \frac{\partial}{\partial \tau} Z^a \right], \quad (4.20)$$

To check that this relation is indeed the field action of a massless particle of Minkowski space, we first replace $\bar{Z}_a$ and Z^a by λ 's and μ 's in $\bar{Z}_a i \frac{\partial}{\partial \tau} Z^a$. This operation leads to $(\bar{\lambda}_{\dot{\alpha}} i \partial_{\tau} \mu^{\dot{\alpha}} - \bar{\mu}_{\alpha} i \partial_{\tau} \lambda^{\alpha})$. Then use incidence relation (4.11) to put the action in the form $\int d\tau (\lambda^{\beta} \bar{\lambda}_{\dot{\alpha}}) (\partial_{\tau} x_{\beta}^{\dot{\alpha}})$ which is nothing but

$$\mathcal{S}_{Minkowski} = \int d\tau (p_{\mu} \partial_{\tau} x^{\mu}), \quad (4.21)$$

the usual Minkowski space action of a free massless particle.

Moreover using eq(4.20) and requiring gauge invariance under local projective symmetry (4.14), the gauge invariant twistor action becomes

$$\mathcal{S}_{twistor} [Z, \bar{Z}, V] = - \int d\tau [\bar{Z}_a i (\partial_{\tau} - iV) Z^a - 2hV] \quad (4.22)$$

where V is a gauge field and h is a Fayet-Iliopoulos (FI) like coupling constant; it is just the helicity of the spinning particle. By eliminating the gauge field V through its equation of motion

$$\frac{\delta \mathcal{S}_{twistor} [V]}{\delta V} = 0, \quad (4.23)$$

we rediscover the classical constraint eq(4.18) $\sum_a \bar{Z}_a Z^a = 2h$ which we prefer to rewrite it as follows,

$$\sum_A (Z^a \bar{Z}_a + \bar{Z}_a Z^a) = 4h \quad (4.24)$$

and from which one recognizes eq(4.18) once h is set to zero.

We end this overview and comments on Penrose twistor gauge theory by saying few words about quantum mechanics. In covariant canonical quantization of above twistor model, the twistor variable $\bar{Z}_a$ (the conjugate momentum of Z^a) gets replaced by $-\frac{\partial}{\partial Z^a}$. As such the twistor space wave functions $\Psi(Z) = \langle Z | \Psi \rangle$ are then holomorphic and homogeneous functions of degree $(-2h - 2)$.

$$\Psi(e^{i\theta} Z) = e^{-i(2h+2)\theta} \Psi(Z). \quad (4.25)$$

This homogeneity feature follows from the quantum promotion of the constraint eq(4.24) which gets then replaced by,

$$Z^a \frac{\partial}{\partial Z^a} \Psi(Z) = (-2h - 2) \Psi(Z) \quad (4.26)$$

We turn now to build our fermionic model.

4.2 Fermionic construction

This pure fermionic construction is important for the derivation of fermionic twistor like theory to be developed in next section. The field model we study here deals with

the dynamics of world line fermions transforming as well in the 4- dimension space time and can be viewed as:

- (i) the fermionic analog of eqs(2.2,4.21) dual to twistor construction à la Penrose.
- (ii) the fermionic sector of a supersymmetric gauge theory along the lines outlined in section 2.

Instead of the world line field bosons $x_{\dot{\beta}}^{\dot{\alpha}}(\tau)$ and $p_{\dot{\alpha}}^{\beta}(\tau)$ considered above, our pure fermionic model involves the following ingredients:

(1) **Degrees of freedom:** The model involves the world line fermions

$$\Upsilon_{-\beta}^{\dot{\alpha}}(\tau), \quad \Upsilon_{+\beta}^{\dot{\alpha}}(\tau), \quad \tilde{\Upsilon}_{-\beta}^{\dot{\alpha}}(\tau), \quad \tilde{\Upsilon}_{+\beta}^{\dot{\alpha}}(\tau) \quad (4.27)$$

(see eqs(2.34) to fix the ideas). These fields carries two kinds of indices: **(a)** An index $(\pm)$ exhibiting the statistics of $\Upsilon_{\pm\beta}^{\dot{\alpha}}$ and $\tilde{\Upsilon}_{\pm\beta}^{\dot{\alpha}}$; they are world line fermions. **(b)** dotted and undotted indices $(\alpha, \dot{\alpha} = 1, 2)$ showing that the fields transform as 4- vectors (bi-spinors) in one of the three possible $\mathbb{R}^{(d,4-d)}$ spaces,

$$\Upsilon_{\pm}^m = (\sigma^m)_{\dot{\beta}}^{\alpha} \Upsilon_{\pm\alpha}^{\dot{\beta}}, \quad \tilde{\Upsilon}_{\pm}^m = (\sigma^m)_{\dot{\beta}}^{\alpha} \tilde{\Upsilon}_{\pm\alpha}^{\dot{\beta}}. \quad (4.28)$$

Though stringy applications suggest to consider $\mathbb{R}^{(1,3)}$, it is also interesting to consider also the $\mathbb{R}^{(2,2)}$ and $\mathbb{R}^4$ cases. Below, we shall mainly focus our attention here on the case $\mathbb{R}^{(2,2)}$ because of specific properties; in particular because of the two following:

- (i) In $\mathbb{R}^{(1,3)}$ and $\mathbb{R}^4$ with rotation groups $SO(1,3, \mathbb{R}) \simeq SL(2, \mathbb{C})$ and $SO(4, \mathbb{R}) \simeq SU(2) \times SU(2)$ respectively, $(\frac{1}{2}, 0)$ and $(0, \frac{1}{2})$ spinors are complex,

$$\tilde{\Upsilon}_{\pm}^m = \overline{(\Upsilon_{\pm}^m)}. \quad (4.29)$$

and the underlying twistor space PT splits into holomorphic and antiholomorphic sectors interchanged under complex conjugation,

$$\left(\frac{1}{2}, 0\right) \longleftrightarrow \left(0, \frac{1}{2}\right) \quad (4.30)$$

This complex structure is rich from algebraic geometry view and stringy applications, in particular in topological string on conifold.

- (ii) In $\mathbb{R}^{(2,2)}$ the rotation group is given by $SO(2,2, \mathbb{R}) \simeq SL(2, \mathbb{R}) \times SL(2, \mathbb{R})$ and basic spinors are real. In this case, it is possible to drop out half of the degrees of freedom by keeping for instance just

$$\Upsilon_{\pm\beta}^{\dot{\alpha}} \sim \Upsilon_{\pm}^m, \quad (4.31)$$

in eqs(4.28).

Note also that since $(\Upsilon_{\pm}^m) = (\Upsilon_{\pm}^0, \Upsilon_{\pm}^1, \Upsilon_{\pm}^2, \Upsilon_{\pm}^4,)$ are fermions, the square of each component $\Upsilon_{\pm}^m$ vanishes individually,

$$(\Upsilon_{\pm}^m)^2 = 0, \quad m = 0, 1, 2, 3. \quad (4.32)$$

Moreover since the metric η_{mn} is diagonal we also have

$$\eta_{mn} \Upsilon_{\pm}^m \Upsilon_{\pm}^n \sim \epsilon^{\alpha\beta} \epsilon_{\dot{\alpha}\dot{\beta}} \Upsilon_{-\alpha}^{\dot{\alpha}} \Upsilon_{+\beta}^{\dot{\beta}} = 0. \quad (4.33)$$

Similar relations may be written down for $\tilde{\Upsilon}_{\pm}^m$.

(2) Field action: The pure fermionic field action $\mathcal{S}_F$ describing the dynamics of the free fields $\Upsilon_{\pm\alpha}^{\dot{\alpha}}$ and $\tilde{\Upsilon}_{-\alpha}^{\dot{\alpha}}$ is given by, see also eq(3.21),

$$\begin{aligned} \mathcal{S}_F [\Upsilon, \tilde{\Upsilon}] &= \int_{\mathbb{R}} d\tau \left[\frac{1}{2} \left(\Upsilon_{-\alpha}^{\dot{\alpha}} \frac{\partial}{\partial \tau} \Upsilon_{+\beta}^{\dot{\beta}} - \Upsilon_{+\beta}^{\dot{\beta}} \frac{\partial}{\partial \tau} \Upsilon_{-\alpha}^{\dot{\alpha}} \right) \epsilon^{\alpha\beta} \epsilon_{\dot{\alpha}\dot{\beta}} \right] \\ &+ \int_{\mathbb{R}} d\tau \left[\frac{1}{2} \left(\tilde{\Upsilon}_{-\alpha}^{\dot{\alpha}} \frac{\partial}{\partial \tau} \tilde{\Upsilon}_{+\beta}^{\dot{\beta}} - \tilde{\Upsilon}_{+\beta}^{\dot{\beta}} \frac{\partial}{\partial \tau} \tilde{\Upsilon}_{-\alpha}^{\dot{\alpha}} \right) \epsilon^{\alpha\beta} \epsilon_{\dot{\alpha}\dot{\beta}} \right]. \end{aligned} \quad (4.34)$$

Note in passing that, under some hypothesis on chirality, the second line may dropped out. Note also that comparing $\mathcal{S}_F$, which can be also written as

$$\begin{aligned} \mathcal{S}_F [\Upsilon_{\pm}, \partial_{\tau} \Upsilon_{\pm}] &\sim \int_{\mathbb{R}} d\tau \left[\eta_{mn} \left(\Upsilon_{-}^m \frac{\partial}{\partial \tau} \Upsilon_{+}^n \right) \right] \\ &+ \int_{\mathbb{R}} d\tau \left[\eta_{mn} \left(\tilde{\Upsilon}_{-}^m \frac{\partial}{\partial \tau} \tilde{\Upsilon}_{+}^n \right) \right], \end{aligned} \quad (4.35)$$

with its Bose analog eq(4.21), one learns the naive correspondence,

$$\begin{aligned} x^n &\longrightarrow \Upsilon_{+}^n, \quad \tilde{\Upsilon}_{+}^n \\ p_m &\longrightarrow \Upsilon_{-m}, \quad \tilde{\Upsilon}_{-m}. \end{aligned} \quad (4.36)$$

(3) The conjugate momentum variables $\pi_{-\dot{\alpha}}^{\beta}$ and $\tilde{\pi}_{-\dot{\alpha}}^{\beta}$,

$$\pi_{-\dot{\alpha}}^{\beta}(\tau) = \frac{\delta \mathcal{S}_F}{\delta (\partial \Upsilon_{+\beta}^{\dot{\alpha}})}, \quad \tilde{\pi}_{-\dot{\alpha}}^{\beta} = \frac{\delta \mathcal{S}_F}{\delta (\partial \tilde{\Upsilon}_{+\beta}^{\dot{\alpha}})}. \quad (4.37)$$

By using the above expression of $\mathcal{S}_F$, we get

$$\pi_{-\dot{\alpha}}^{\alpha} = -\epsilon^{\alpha\beta} \epsilon_{\dot{\alpha}\dot{\beta}} \Upsilon_{-\beta}^{\dot{\beta}}, \quad \tilde{\pi}_{-\dot{\alpha}}^{\alpha} = -\epsilon^{\alpha\beta} \epsilon_{\dot{\alpha}\dot{\beta}} \tilde{\Upsilon}_{-\beta}^{\dot{\beta}}. \quad (4.38)$$

This relation shows that the fermionic fields $\Upsilon_{+\beta}^{\dot{\beta}}$, $\tilde{\Upsilon}_{+\beta}^{\dot{\beta}}$ and $\Upsilon_{-\beta}^{\dot{\beta}}$, $\tilde{\Upsilon}_{-\beta}^{\dot{\beta}}$ are conjugate to each other. So to avoid confusions, it is interesting to re-parameterize the above fermionic field action as follows:

$$\mathcal{S}_F [\Upsilon, \pi, \tilde{\Upsilon}, \tilde{\pi}] = \int_{\mathbb{R}} d\tau \text{Tr} \left(\pi_{-\dot{\alpha}}^{\alpha} \frac{\partial}{\partial \tau} \Upsilon_{+\alpha}^{\dot{\alpha}} + \tilde{\pi}_{-\dot{\alpha}}^{\alpha} \frac{\partial}{\partial \tau} \tilde{\Upsilon}_{+\alpha}^{\dot{\alpha}} \right) \quad (4.39)$$

Note also because of the fermionic nature of the $\pi_{-\dot{\alpha}}^{\alpha}$'s and like for eq(4.33), we have here also the nilpotency property

$$\text{Tr} (\pi_{-}^2) = \pi_{-\dot{\alpha}}^{\alpha} \pi_{-\alpha}^{\dot{\alpha}} = 0, \quad \text{Tr} (\tilde{\pi}_{-}^2) = \tilde{\pi}_{-\dot{\alpha}}^{\alpha} \tilde{\pi}_{-\alpha}^{\dot{\alpha}} = 0. \quad (4.40)$$

Since the pairs $(\Upsilon_{+\beta}^{\dot{\beta}}, \pi_{-\dot{\alpha}}^{\alpha})$ and $(\tilde{\Upsilon}_{+\dot{\beta}}^{\beta}, \tilde{\pi}_{-\dot{\alpha}}^{\alpha})$ play quite similar role, we shall fix our attention on one of them, say $(\Upsilon_{+\beta}^{\dot{\beta}}, \pi_{-\dot{\alpha}}^{\alpha})$ with field action

$$\mathcal{S}_F[\Upsilon, \pi] = \int_{\mathbb{R}} d\tau \text{Tr} \left(\pi_{-\dot{\alpha}}^{\alpha} \frac{\partial}{\partial \tau} \Upsilon_{+\alpha}^{\dot{\alpha}} \right), \quad (4.41)$$

then give the results for $(\tilde{\Upsilon}_{+\dot{\beta}}^{\beta}, \tilde{\pi}_{-\dot{\alpha}}^{\alpha})$ whenever needed.

4.2.1 Basic symmetries

In the case of $\mathbb{R}^{(2,2)}$ geometry, the field action $\mathcal{S}_F$ has the following remarkable and manifest symmetries:

- (1) $SO(2, 2, \mathbb{R}) \sim SL(2, \mathbb{R}) \times SL(2, \mathbb{R})$ global invariance.
- (2) Invariance under the real scaling transformation

$$\begin{aligned} \Upsilon_{+\alpha}^{\dot{\alpha}} &\longrightarrow e^{+v} \Upsilon_{+\alpha}^{\dot{\alpha}} \\ \pi_{-\dot{\alpha}}^{\alpha} &\longrightarrow e^{-v} \pi_{-\dot{\alpha}}^{\alpha}. \end{aligned} \quad (4.42)$$

with a global real parameter $v \in \mathbb{R}$. It captures the spin of the world line fields and has also a world sheet interpretation in string theory.

Moreover promoting this symmetry to a local one, that is $v = v(\tau)$ or equivalently,

$$\frac{\partial v}{\partial \tau} \neq 0, \quad (4.43)$$

then gauge invariance requires that the above field action (4.41) should be extended as,

$$\mathcal{S}_{FG}[\Upsilon_+, \pi_-, \mathcal{A}] \sim \int d\tau \text{Tr} \left[\pi_{-\dot{\alpha}}^{\alpha} \left(\frac{\partial}{\partial \tau} - \mathcal{A}_{\tau} T_0 \right) \Upsilon_{+\alpha}^{\dot{\alpha}} \right] \quad (4.44)$$

where $\mathcal{A} = \mathcal{A}(\tau)$ is the gauge field associated with the above real scaling invariance. In this relation, T_0 is the generator of the scaling symmetry (4.42) and acts on the fields Υ_+ and π_- like:

$$[T_0, \Upsilon_{+\alpha}^{\dot{\alpha}}] = +\frac{1}{2} \Upsilon_{+\alpha}^{\dot{\alpha}}, \quad [T_0, \pi_{-\dot{\alpha}}^{\alpha}] = -\frac{1}{2} \pi_{-\dot{\alpha}}^{\alpha}. \quad (4.45)$$

The meaning of this gauge invariance can be made transparent by noting that the gauge field $\mathcal{A}_{\tau}$ do not propagate and so it is an auxiliary field. Eliminating it through its equation of motion

$$\frac{\delta \mathcal{S}_{FG}}{\delta \mathcal{A}_{\tau}} = 0, \quad (4.46)$$

we get the following constraint eq,

$$\text{Tr} (\pi_{-\dot{\alpha}}^{\alpha} \Upsilon_{+\alpha}^{\dot{\alpha}}) = 0. \quad (4.47)$$

But this equation is a familiar QFT relation; it captures the antisymmetry property of Fermi like fields as it can be seen by using the cyclic property of trace, $\text{Tr} (\pi_{-\dot{\alpha}}^{\alpha} \Upsilon_{+\alpha}^{\dot{\alpha}} + \Upsilon_{+\alpha}^{\dot{\alpha}} \pi_{-\dot{\alpha}}^{\alpha}) = 0$.

Theorem 1

In pure fermionic phase space with canonical coordinate fields $\Upsilon_{+\alpha}^{\dot{\alpha}}$ ($\tilde{\Upsilon}_{+\alpha}^{\dot{\alpha}}$) and conjugate momenta $\pi_{-\dot{\alpha}}^{\alpha}$ ($\tilde{\pi}_{-\dot{\alpha}}^{\alpha}$), the free field action $\mathcal{S}_{FG}$ ($\tilde{\mathcal{S}}_{FG}$) has a local $SP(2, \mathbb{R})$ gauge invariance capturing the three Fermi field antisymmetry properties

$$\begin{aligned}\text{Tr}(\pi_{-\dot{\alpha}}^{\alpha} \Upsilon_{+\alpha}^{\dot{\alpha}}) &= 0 \\ \text{Tr}(\pi_{-\dot{\alpha}}^{\alpha} \pi_{-\alpha}^{\dot{\alpha}}) &= 0 \\ \text{Tr}(\Upsilon_{+\dot{\alpha}}^{\alpha} \Upsilon_{+\alpha}^{\dot{\alpha}}) &= 0.\end{aligned}\tag{4.48}$$

These are just the constraint relations derived above; see eqs(4.33,4.40,4.47). Similar relations are also valid for $\tilde{\Upsilon}_{+\dot{\beta}}^{\beta}$ and $\tilde{\pi}_{-\dot{\alpha}}^{\alpha}$.

4.2.2 More on $SP(2, \mathbb{R})$ gauge invariance

Before giving the $SP(2, \mathbb{R})$ gauge invariant field action, we begin by recalling some useful tools on the real group $SP(2, \mathbb{R})$ and some aspects of its representations. Its Lie algebra $sp(2, \mathbb{R})$ is three dimensional with generators T_{--}, T_0 and T_{++} satisfying the commutation relations,

$$\begin{aligned}[T_{--}, T_{++}] &= 2T_0, \\ [T_0, T_{++}] &= T_{++}, \\ [T_0, T_{--}] &= -T_{--},\end{aligned}\tag{4.49}$$

where $\Delta = 2T_0$ is the operator counting the charges of $SP(2, \mathbb{R})$ field representations. In particular we have,

$$[\Delta, T_{\pm\pm}] = \pm 2T_{\pm\pm}, \quad [\Delta, \Upsilon_{+\alpha}^{\dot{\alpha}}] = +\Upsilon_{+\alpha}^{\dot{\alpha}}, \quad [\Delta, \pi_{-\dot{\alpha}}^{\alpha}] = -\pi_{-\dot{\alpha}}^{\alpha}.\tag{4.50}$$

There are different ways to realize this real symmetry; one of these ways is given by the following left moving wold sheet differential operators,

$$T_{--} = \frac{\partial}{\partial \sigma^+}, \quad T_0 = \sigma^+ \frac{\partial}{\partial \sigma^+}, \quad T_{++} = \sigma^{+2} \frac{\partial}{\partial \sigma^+}.\tag{4.51}$$

from which one can read immediately their scaling behaviours under global the transformations $\sigma^+ \longrightarrow e^{2v} \sigma^+$,

$$T'_{--} = e^{-2v} T_{--}, \quad T'_0 = T_0, \quad T'_{++} = e^{2v} T_{++}.\tag{4.52}$$

The $T_{0,\pm\pm}$'s can be also realized as 2×2 matrices when acting on two component vectors. This is the case which interests us here. The 2-vector is given by (Υ_+, π_-) and the $T_{0,\pm\pm}$

matrices act as,

$$\begin{aligned}
[T_{++}, \Upsilon_+] &= 0, & [T_{--}, \pi_-] &= 0 \\
[T_{--}, \Upsilon_+] &\sim \pi_-, & [T_{++}, \pi_-] &\sim \Upsilon_+ \\
[T_0, \Upsilon_+] &= +\frac{1}{2}\Upsilon_+, & [T_0, \pi_-] &= -\frac{1}{2}\pi_-
\end{aligned} \tag{4.53}$$

We come now to derive the general expression of the classical field action $\mathcal{S}_{FG}[\Upsilon_+, \pi_-, \mathcal{B}]$ extending eq(4.44) and recovering the constraint eqs(4.48). This action has an $SP(2, \mathbb{R})$ gauge invariance and reads as,

$$\mathcal{S}_{FG}[\Upsilon_+, \pi_-, \mathcal{B}] \sim \int d\tau \left(\text{Tr} \left[\pi_{-\dot{\alpha}}^{\alpha} \left(\frac{\partial}{\partial \tau} - \mathcal{B}_{\tau} \right) \Upsilon_{+\alpha}^{\dot{\alpha}} \right] \right) \tag{4.54}$$

where now the gauge field $\mathcal{B}_{\tau}$ is valued in the $sp(2, \mathbb{R})$ algebra, i.e,

$$\mathcal{B}_{\tau} = \mathcal{B}_{\tau,0}T_0 + \mathcal{B}_{\tau,-}T_{++} + \mathcal{B}_{\tau,+}T_{--}. \tag{4.55}$$

Eq(4.54) can be put into a more condensed form by using the $SP(2, \mathbb{R})$ spinor $\Gamma^i = \epsilon^{ij}\Gamma_j$ with $i = +, -$, $\epsilon^{+-} = 1$; that is

$$\Gamma_i = (\pi_-, -\Upsilon_+), \quad \Gamma^i = \begin{pmatrix} \Upsilon_+ \\ \pi_- \end{pmatrix}, \tag{4.56}$$

We have

$$\mathcal{S}_{FG}[\Gamma, \mathcal{B}] \sim \int d\tau \left(\frac{1}{2} \text{Tr} \left[\Gamma_{\dot{\alpha}}^{i\alpha} \left(\epsilon_{ij} \frac{\partial}{\partial \tau} - (\mathcal{B}_{\tau})_{(ij)} \right) \Gamma_{\alpha}^{j\dot{\alpha}} \right] \right) \tag{4.57}$$

From which one can re-express the constraint eqs(4.48) using the world line Γ fields. Similar results are valid for the twild sector $\tilde{\Gamma}$ and $\tilde{\mathcal{B}}$.

5 Twistor like dual

To build the twistor representation dual to the above the pure fermionic model, we proceed as in bosonic case. Below, we summarize the steps of the procedure:

5.1 Solve the constraint eqs

First solve the constraint eqs(4.48) by using fermionic twistor like variables. Following the idea outlined in introduction and using preliminary results of section 2; then mimicking the usual bosonic twistor analysis, it is not difficult to see that the solution of the constraint eqs is given by,

$$\begin{aligned}
\Upsilon_{+\alpha}^{\dot{\alpha}} &= \lambda_{\alpha} \zeta_{+}^{\dot{\alpha}}, & \tilde{\Upsilon}_{+\alpha}^{\dot{\alpha}} &= \zeta_{+\alpha} \lambda^{\dot{\alpha}} \\
\pi_{-\dot{\alpha}}^{\alpha} &= \lambda^{\alpha} \zeta_{-\dot{\alpha}}, & \tilde{\pi}_{-\dot{\alpha}}^{\alpha} &= \zeta_{-\alpha} \lambda^{\dot{\alpha}},
\end{aligned} \tag{5.1}$$

where, for simplicity, we have set dropped out the hats on the fields; $\tilde{\zeta}_{\pm}^{\dot{\alpha}} \equiv \zeta_{\pm}^{\dot{\alpha}}$ and $\tilde{\lambda}^{\dot{\alpha}} \equiv \lambda^{\dot{\alpha}}$. In this solution, the λ_{α} 's ($\lambda^{\dot{\alpha}}$'s) are commuting world line bosons with the usual property $(\lambda_{\alpha}\lambda^{\alpha}) = 0$; the same λ_{α} 's as in section 3. The $\zeta_{\pm}^{\dot{\alpha}}$'s ($\zeta_{\pm\alpha}$'s) are anticommuting world line fermions capturing the Fermi statistics of $\Upsilon_{+\alpha}^{\dot{\alpha}}$ and $\pi_{-\dot{\alpha}}^{\alpha}$ ($\tilde{\Upsilon}_{+\alpha}^{\dot{\alpha}}$ and $\tilde{\pi}_{-\dot{\alpha}}^{\alpha}$). By substituting the above solution in the anticommutator,

$$\left\{ \Upsilon_{+\alpha}^{\dot{\alpha}}, \pi_{-\dot{\beta}}^{\beta} \right\} = 0, \quad (5.2)$$

we get in a first stage $\left\{ \lambda_{\alpha} \zeta_{+}^{\dot{\alpha}}, \lambda^{\beta} \zeta_{-\dot{\beta}} \right\} = \lambda_{\alpha} \lambda^{\beta} \left\{ \zeta_{+}^{\dot{\alpha}}, \zeta_{-\dot{\beta}} \right\}$ and so the following identities,

$$\left\{ \zeta_{-\dot{\alpha}}, \zeta_{+}^{\dot{\beta}} \right\} = 0, \quad [\lambda_{\alpha}, \lambda^{\beta}] = 0, \quad \left[\lambda_{\alpha}, \zeta_{\pm}^{\dot{\beta}} \right] = 0, \quad (5.3)$$

showing that $\Upsilon_{+\alpha}^{\dot{\alpha}}$ is built out of the product of boson λ_{α} with positive $SO(2, 2, R)$ chirality and a world line fermion with negative $SO(2, 2, R)$ chirality. The opposite combination corresponds exactly to $\pi_{-\dot{\alpha}}^{\alpha}$.

Moreover, notice that the solution (5.1) is not unique since arbitrary changes of the $SO(2, 2, R)$ spinors of the form

$$\begin{aligned} \lambda^{\alpha} &\rightarrow \lambda^{\alpha'} = e^{\varphi} \lambda^{\alpha}, \\ \zeta_{\pm}^{\dot{\alpha}} &\rightarrow \zeta_{\pm}^{\dot{\alpha}'} = e^{-\varphi} \zeta_{\pm}^{\dot{\alpha}}, \end{aligned} \quad (5.4)$$

with $\varphi = \varphi(\tau)$ is an arbitrary real number; $\varphi \in \mathbb{R}$, leaves eqs(5.1) invariant. This transformation corresponds to the usual complex projective symmetry of bosonic twistor space. Notice also that under the scaling transformations $\Upsilon_{+\alpha}^{\dot{\alpha}} \rightarrow e^v \Upsilon_{+\alpha}^{\dot{\alpha}}$ and $\pi_{-\dot{\alpha}}^{\alpha} \rightarrow e^{-v} \pi_{-\dot{\alpha}}^{\alpha}$ eqs(5.1), we have moreover,

$$\lambda^{\alpha} \rightarrow \lambda^{\alpha}, \quad \zeta_{\pm}^{\dot{\alpha}} \rightarrow e^{\pm v} \zeta_{\pm}^{\dot{\alpha}}, \quad (5.5)$$

and should not be confused with eq(5.4). As this symmetry captures just the Fermi statistics of the $\zeta_{\pm}$'s, we shall forget it and then focus our attention in what follows on the real projective invariance (5.4). This symmetry plays a crucial role in the derivation fermionic twistor like model and its quantization.

5.2 Twistor field action $\mathcal{S}_{FT}$

To build the fermionic twistor like field action $\mathcal{S}_{FT}$ dual to eq(4.34) and derive the underlying gauge constraint eqs, we start by introducing the projective twistor variables

$$z^a, w_a, \quad (5.6)$$

for the fermionic twistor like theory. Next we derive the field action $\mathcal{S}_{FT} = \mathcal{S}_{FT}[z^a, w_a]$ of the classical model and consider then its quantization.

Roughly, twistor variables are twistor phase space coordinates, given in the case of a massless particle moving in $\mathbb{R}^{(1,3)}$, by

$$\lambda^\alpha, \mu^{\dot{\alpha}}; \bar{\mu}_\alpha, \bar{\lambda}_{\dot{\alpha}}, \quad (5.7)$$

where $\mu^{\dot{\alpha}} = \frac{\delta S_{BT}}{\delta \partial_\tau \lambda_{\dot{\alpha}}}$ and $\bar{\mu}_\alpha = \frac{\delta S_{BT}}{\delta \partial_\tau \lambda^\alpha}$. It happens that these twistor variables split into complex homogeneous holomorphic variables

$$z^a = (\lambda^\alpha, \mu^{\dot{\alpha}}) \in CP^3, \quad (5.8)$$

and their complex antiholomorphic ones $\bar{z}_a = (\bar{\lambda}_{\dot{\alpha}}, \bar{\mu}_\alpha)$. In the case of a massless particle moving in the $\mathbb{R}^{(2,2)}$ geometry, the analysis is a little bit different for the two following:

(i) Spinors of $SO(2, 2, \mathbb{R})$ target space group are real; so the corresponding twistor space is real and apparently there is no holomorphic and anti-holomorphic like splitting as in the complex case. However, we do still have the two sectors given by $(\Upsilon_{+\alpha}^{\dot{\alpha}}, \pi_{-\dot{\alpha}}^\alpha)$ and $(\tilde{\Upsilon}_{+\alpha}^{\dot{\alpha}}, \tilde{\pi}_{-\dot{\alpha}}^\alpha)$, but stable under complex conjugation. More precisely, we have for the $(\Upsilon_{+\alpha}^{\dot{\alpha}}, \pi_{-\dot{\alpha}}^\alpha)$ sector the twistor like variables,

$$(\lambda^1, \lambda^2, \xi_{-1}, \xi_{-2}, \nu_1, \nu_2, \zeta_+^1, \zeta_+^2) \in WRP_{(+1,+1,+1,+1,-1,-1,-1,-1)}^{(3|4)}, \quad (5.9)$$

where $(+1, +1, +1, +1, -1, -1, -1, -1)$ stand for the weight vector of the scaling transformation of the super multiplet $(\lambda^\alpha, \xi_{-\dot{\alpha}}, \nu_\alpha, \zeta_+^{\dot{\alpha}})$. The space $WRP_{(++++----)}^{(3|4)}$ stands for the weighted real projective supermanifold. For $(\tilde{\Upsilon}_{+\alpha}^{\dot{\alpha}}, \tilde{\pi}_{-\dot{\alpha}}^\alpha)$ sector, we have the following projective supercoordinates,

$$(\zeta_+^\alpha, \nu^{\dot{\alpha}}, \xi_{-\alpha}, \lambda_{\dot{\alpha}}). \quad (5.10)$$

In first sector bosons λ^α and ν_α are in $(\frac{1}{2}, 0)$ representation of $SO(2, 2, \mathbb{R})$ while in the second, $\lambda_{\dot{\alpha}}$ and $\nu^{\dot{\alpha}}$ are in $(0, \frac{1}{2})$. Fermions have the opposite chirality.

(ii) The real field variables

$$(\lambda^\alpha, \lambda_{\dot{\alpha}} | \zeta_+^\alpha, \zeta_{+\dot{\alpha}}), \quad (5.11)$$

and their conjugate momenta,

$$(\nu_\alpha, \nu^{\dot{\alpha}} | \xi_{-\alpha}, \xi_{-\dot{\alpha}}). \quad (5.12)$$

are related as,

$$\begin{aligned} \nu_\alpha &= \frac{\delta \mathcal{S}_{FT}}{\delta (\partial_\tau \lambda^\alpha)}, & \nu^{\dot{\alpha}} &= \frac{\delta \mathcal{S}_{FT}}{\delta (\partial_\tau \lambda_{\dot{\alpha}})}, \\ \xi_{-\alpha} &= \frac{\delta \mathcal{S}_{FT}}{\delta (\partial_\tau \zeta_+^\alpha)}, & \xi_{-\dot{\alpha}} &= \frac{\delta \mathcal{S}_{FT}}{\delta (\partial_\tau \zeta_{+\dot{\alpha}})}. \end{aligned} \quad (5.13)$$

and parameterize the full phase space of the fermionic twistor like model,

$$\mathcal{E}_{phase} = \{ \lambda^\alpha, \lambda_{+\dot{\alpha}}, \nu_\alpha, \nu^{\dot{\alpha}} | \zeta_+^\alpha, \zeta_+^{\dot{\alpha}}, \xi_{-\alpha}, \xi_{-\dot{\alpha}} \}, \quad (5.14)$$

These variables satisfy the following graded canonical relations,

$$\begin{aligned}\{\lambda^\alpha, \nu_\beta\}_{GPB} &= \delta^\alpha_\beta, & \{\lambda_{+\dot{\beta}}, \nu^{\dot{\alpha}}\}_{GPB} &= \delta^{\dot{\alpha}}_{\dot{\beta}} \\ \{\zeta_+^\alpha, \xi_{-\beta}\}_{GPB} &= \delta^\alpha_\beta, & \{\zeta_+^{\dot{\alpha}}, \xi_{-\dot{\beta}}\}_{GPB} &= \delta^{\dot{\alpha}}_{\dot{\beta}},\end{aligned}\quad (5.15)$$

where the graded Poisson bracket $\{f, g\}_{GPB}$, of generic functions $f = f(\lambda, \nu | \zeta_+, \xi_-)$ and $g = g(\lambda, \nu | \zeta_+, \xi_-)$ living on twistor space, is defined as

$$\begin{aligned}\{f, g\}_{GPB} &= \left(\frac{\partial f}{\partial \lambda^\alpha} \frac{\partial g}{\partial \nu_\alpha} - \frac{\partial f}{\partial \nu_\alpha} \frac{\partial g}{\partial \lambda^\alpha} \right) + \left(\frac{\partial f}{\partial \zeta_+^\alpha} \frac{\partial g}{\partial \xi_{-\alpha}} + \frac{\partial f}{\partial \xi_{-\alpha}} \frac{\partial g}{\partial \zeta_+^\alpha} \right) \\ &+ \left(\frac{\partial f}{\partial \lambda_{+\dot{\alpha}}} \frac{\partial g}{\partial \nu^{\dot{\alpha}}} - \frac{\partial f}{\partial \nu^{\dot{\alpha}}} \frac{\partial g}{\partial \lambda_{+\dot{\alpha}}} \right) + \left(\frac{\partial f}{\partial \zeta_+^{\dot{\alpha}}} \frac{\partial g}{\partial \xi_{-\dot{\alpha}}} + \frac{\partial f}{\partial \xi_{-\dot{\alpha}}} \frac{\partial g}{\partial \zeta_+^{\dot{\alpha}}} \right).\end{aligned}\quad (5.16)$$

For convenience and later use, it is interesting to combine these variables into super-coordinates as,

$$\begin{aligned}z^a &= (\lambda^\alpha | \xi_{-\dot{\alpha}}), & \tilde{z}^a &= (\zeta_+^\alpha | \nu^{\dot{\alpha}}), \\ w^b &= (\nu^\beta | \zeta_+^{\dot{\beta}}), & \tilde{w}^b &= (\xi_{-\dot{\beta}}^\beta | \lambda^{\dot{\beta}}).\end{aligned}\quad (5.17)$$

With these super variables, the real projective transformations of the twistor superspace,

$$\begin{aligned}\lambda^\alpha &\rightarrow e^{+\varphi} \lambda^\alpha, & \zeta_+^\alpha &\rightarrow e^{+\theta} \zeta_+^\alpha, \\ \nu_\alpha &\rightarrow e^{-\varphi} \nu_\alpha, & \xi_{-\alpha} &\rightarrow e^{-\theta} \xi_{-\alpha}, \\ \zeta_+^{\dot{\alpha}} &\rightarrow e^{-\varphi} \zeta_+^{\dot{\alpha}}, & \lambda_{\dot{\alpha}} &\rightarrow e^{-\theta} \lambda_{\dot{\alpha}}, \\ \xi_{-\dot{\alpha}} &\rightarrow e^{+\varphi} \xi_{-\dot{\alpha}}, & \nu^{\dot{\alpha}} &\rightarrow e^{+\theta} \nu^{\dot{\alpha}},\end{aligned}\quad (5.18)$$

combine as follows,

$$\begin{aligned}z^a &\rightarrow e^{+\varphi} z^a, & \tilde{z}^a &\rightarrow e^{+\theta} \tilde{z}^a, \\ w^a &\rightarrow e^{-\varphi} w^a, & \tilde{w}^a &\rightarrow e^{-\theta} \tilde{w}^a.\end{aligned}\quad (5.19)$$

Note that the parameters φ and θ are a priori independent, unless if the momentum vector,

$$p_\alpha^\alpha = \lambda^\alpha \tilde{\lambda}_{\dot{\alpha}} \quad (5.20)$$

is required to be invariant under (5.18). In this case, we should have the identity,

$$\varphi = \theta. \quad (5.21)$$

Note also that within the $(\Upsilon_{+\alpha}^{\dot{\alpha}}, \pi_{-\dot{\alpha}}^\alpha)$ sector eq(5.9), the twistor like variables $(\lambda^\alpha, \xi_{-\dot{\alpha}}, \nu_\alpha, \zeta_+^{\dot{\alpha}})$ splits into blocs as follows,

$$\begin{aligned}z^a &= (\lambda^\alpha | \xi_{-\dot{\alpha}}) \in RP_{(+1, +1, +1, +1)}^{(1|2)}, \\ w^b &= (\nu^\beta | \zeta_+^{\dot{\beta}}) \in RP_{(-1, -1, -1, -1)}^{(1|2)}.\end{aligned}\quad (5.22)$$

Moreover using the identities (5.13), the component twistor field action reads, up to gauge coupling terms type $\xi_{\mp\dot{\alpha}} V \zeta_{\pm}^{\dot{\alpha}}$ to be given later, as follows,

$$\begin{aligned} \mathcal{S}_{FT} = & \frac{c_1}{2} \int d\tau [\nu_{\alpha} \partial_{\tau} \lambda^{\alpha} - \lambda^{\alpha} (\partial_{\tau} \nu_{\alpha})] \\ & - \frac{c'_1}{2} \int d\tau [\nu^{\dot{\alpha}} \partial_{\tau} \lambda_{\dot{\alpha}} - \lambda_{\dot{\alpha}} (\partial_{\tau} \nu^{\dot{\alpha}})] \\ & - \frac{c_2}{2} \int d\tau [\xi_{-}^{\dot{\alpha}} \partial_{\tau} \zeta_{+\dot{\alpha}} + \zeta_{+\dot{\alpha}} (\partial_{\tau} \xi_{-}^{\dot{\alpha}})] \\ & + \frac{c'_2}{2} \int d\tau [\xi_{-\alpha} \partial_{\tau} \zeta_{+}^{\alpha} + \zeta_{+}^{\alpha} (\partial_{\tau} \xi_{-\alpha})]. \end{aligned} \quad (5.23)$$

The number c_1 , c'_1 , c_2 and c'_2 are determined by requiring this action to coincide with the action (4.39),

$$\mathcal{S}_{FT} [\Upsilon, \pi, \tilde{\Upsilon}, \tilde{\pi}] = \int_{\mathbb{R}} d\tau \text{Tr} \left(\pi_{-\dot{\alpha}}^{\alpha} \frac{\partial}{\partial \tau} \Upsilon_{+\alpha}^{\dot{\alpha}} + \tilde{\pi}_{-\dot{\alpha}}^{\alpha} \frac{\partial}{\partial \tau} \tilde{\Upsilon}_{+\alpha}^{\dot{\alpha}} \right). \quad (5.24)$$

This requirement allows also to determine the fermionic analog of the Penrose incidence relation (4.11). Setting,

$$\begin{aligned} \nu_{\alpha} &= \pi_{-\alpha}^{\dot{\alpha}} \zeta_{+\dot{\alpha}}, \\ \xi_{-\dot{\alpha}} &= \pi_{-\dot{\alpha}}^{\alpha} \lambda_{\alpha}, \end{aligned} \quad (5.25)$$

and similarly

$$\begin{aligned} \xi_{-}^{\alpha} &= \tilde{\pi}_{-\dot{\alpha}}^{\alpha} \lambda^{\dot{\alpha}}, \\ \nu^{\dot{\alpha}} &= \tilde{\pi}_{-\alpha}^{\dot{\alpha}} \zeta_{+}^{\alpha}. \end{aligned} \quad (5.26)$$

Then putting back into eq(5.23), we get for the integrand $(\nu_{\alpha} \partial_{\tau} \lambda^{\alpha} + c_1 \xi_{-\dot{\alpha}} \partial_{\tau} \zeta_{+}^{\dot{\alpha}})$ the following,

$$\pi_{-\dot{\alpha}}^{\alpha} \zeta_{+}^{\dot{\alpha}} \partial_{\tau} \lambda_{\alpha} + \pi_{-\dot{\alpha}}^{\alpha} \lambda_{\alpha} \partial_{\tau} \zeta_{+}^{\dot{\alpha}} \quad (5.27)$$

provided

$$c_1 = c_2 = 1. \quad (5.28)$$

A similar result is valid for the twild sector. Moreover using the supervariables z^a and w^b , one can rewrite the previous twistor field action as follows,

$$\begin{aligned} \mathcal{S}_{FT} [z, w, \tilde{z}, \tilde{w}] = & \frac{1}{2} \int d\tau [w^b \partial_{\tau} z^a - (\partial_{\tau} w^b) z^a] \omega_{ab} \\ & + \frac{1}{2} \int d\tau [\tilde{w}^b \partial_{\tau} \tilde{z}^a - (\partial_{\tau} \tilde{w}^b) \tilde{z}^a] \omega_{ab} \end{aligned} \quad (5.29)$$

with metric ω_{ab} given by

$$\omega_{ab} = \begin{pmatrix} \epsilon_{\alpha\beta} & 0 \\ 0 & \epsilon_{\dot{\alpha}\dot{\beta}} \end{pmatrix}, \quad \omega^{ba} = \begin{pmatrix} \epsilon^{\beta\alpha} & 0 \\ 0 & \epsilon^{\dot{\beta}\dot{\alpha}} \end{pmatrix}. \quad (5.30)$$

From these eqs, one reads a set of features; in particular the following:

(1) the variable $w_a = \omega_{ab} w^b$ is the conjugate momentum of z^a and so we have amongst others,

$$\begin{aligned}\{z^a, w^b\}_{GPB} &= \omega^{ba}, \\ \{z^a, z^b\}_{GPB} &= \{w_a, w_b\}_{GPB} = 0,\end{aligned}\tag{5.31}$$

where ω^{ab} stands for the inverse of ω_{ba} ; that is $\omega^{ab}\omega_{bc} = \delta_c^a$. We also have for generic functions $f = f(z, w, \tilde{z}, \tilde{w})$ and $g = g(z, w, \tilde{z}, \tilde{w})$, the following

$$\begin{aligned}\{f, g\}_{GPB} &= \omega^{ba} \left(\frac{\partial f}{\partial z^a} \frac{\partial g}{\partial w^b} - (-)^{|a| \cdot |b|} \frac{\partial f}{\partial w^b} \frac{\partial g}{\partial z^a} \right) \\ &+ \omega^{ba} \left(\frac{\partial f}{\partial \tilde{z}^a} \frac{\partial g}{\partial \tilde{w}^b} - (-)^{|a| \cdot |b|} \frac{\partial f}{\partial \tilde{w}^b} \frac{\partial g}{\partial \tilde{z}^a} \right),\end{aligned}\tag{5.32}$$

where $|a| \equiv \text{grad}(a)$ with,

$$\text{grad}(a) = \begin{cases} \text{grad}(\alpha) = 0 \\ \text{grad}(\pm\dot{\alpha}) = 1 \end{cases}\tag{5.33}$$

(2) Eq(5.29) is manifestly invariant under global projective transformations (5.19) and $SL(2|2, R)$ supergroup of the twistor supermanifold $RP^{(1|2)}$.

5.3 Quantum constraints

Promoting the projective transformations (5.18-5.19) to local ones $\varphi = \varphi(\tau)$; $\theta = \theta(\tau)$ that is $\partial_\tau \varphi \neq 0$, $\partial_\tau \theta \neq 0$, then gauge invariance of the field action (5.29) requires implementation of real gauge fields,

$$V_\tau(\tau) \equiv V(\tau), \quad \tilde{V}_\tau(\tau) \equiv \tilde{V}(\tau)\tag{5.34}$$

with gauge transformations

$$\begin{aligned}V' &= V + \partial_\tau \varphi, \\ \tilde{V}' &= \tilde{V} + \partial_\tau \theta.\end{aligned}$$

Thus, the action (5.29) extends as,

$$\begin{aligned}\mathcal{S}_{FT} [z, w, V, \tilde{z}, \tilde{w}, \tilde{V}] &= \frac{1}{2} \left(\int d\tau [w^b D_\tau z^a - (D_\tau w^b) z^a] \omega_{ab} \right) \\ &+ \frac{1}{2} \left(\int d\tau [\tilde{w}^b \tilde{D}_\tau \tilde{z}^a - (\tilde{D}_\tau \tilde{w}^b) \tilde{z}^a] \omega_{ab} \right) \\ &+ \int d\tau 2hV + \int d\tau 2\tilde{h}\tilde{V}\end{aligned}\tag{5.35}$$

where the gauge covariant derivatives are given by,

$$D_\tau z^a = (\partial_\tau - V) z^a, \quad D_\tau w^b = (\partial_\tau + V) w^b \quad (5.36)$$

and

$$\tilde{D}_\tau \tilde{z}^a = (\partial_\tau - \tilde{V}) \tilde{z}^a, \quad \tilde{D}_\tau \tilde{w}^b = (\partial_\tau + \tilde{V}) \tilde{w}^b \quad (5.37)$$

The constants h and $\tilde{h}$ are a priori real coupling constants; they take discrete values at quantum level. Since V and $\tilde{V}$ are non propagating fields, their elimination by using the equations of motion

$$\frac{\delta \mathcal{S}_{FT}}{\delta V} = 0, \quad \frac{\delta \mathcal{S}_{FT}}{\delta \tilde{V}} = 0$$

and lead to the following constraint eqs,

$$\begin{aligned} \omega_{ab} w^b z^a - 2h &= 0 \\ \omega_{ab} \tilde{w}^b \tilde{z}^a - 2\tilde{h} &= 0 \end{aligned} \quad (5.38)$$

or equivalently by using (5.17),

$$\begin{aligned} (\nu_\alpha \lambda^\alpha - \zeta_{+\dot{\alpha}} \xi_-^{\dot{\alpha}}) - 2h &= 0 \\ (\nu_{\dot{\alpha}} \lambda^{\dot{\alpha}} - \zeta_{+\alpha} \xi_-^\alpha) - 2\tilde{h} &= 0 \end{aligned} \quad (5.39)$$

Note that by substituting ν_α and $\xi_-^{\dot{\alpha}}$ and similarly $\nu_{\dot{\alpha}}$ and ξ_-^α by the relations (5.25) associated with helicity zero particles ($h = 0, \tilde{h} = 0$), one finds that $(\nu_\alpha \lambda^\alpha - \zeta_{+\dot{\alpha}} \xi_-^{\dot{\alpha}})$ and $(\nu_{\dot{\alpha}} \lambda^{\dot{\alpha}} - \zeta_{+\alpha} \xi_-^\alpha)$ vanish identically. Note also that eq(5.38) can be also rewritten as,

$$\begin{aligned} \omega_{ab} \left(w^b z^a + (-)^{|a| \cdot |b|} z^a w^b \right) - 4h &= 0, \\ \omega_{ab} \left(\tilde{w}^b \tilde{z}^a + (-)^{|a| \cdot |b|} \tilde{z}^a \tilde{w}^b \right) - 4\tilde{h} &= 0 \end{aligned} \quad (5.40)$$

Quantum mechanically, the graded bracket $\{z^a, w^b\}_{GPB} = \omega^{ba}$ of classical twistor model gets replaced by a graded commutator,

$$[Z^a, W^b] = Z^a W^b + (-)^{|a| \cdot |b|} W^b Z^a = \omega^{ba}. \quad (5.41)$$

A similar relation is valid for the twild variables. Eq(5.41) and its twild analog may be solved like,

$$\begin{aligned} Z^a &= z^a, & W^b &= (-)^{|a| \cdot |b|} \omega^{ba} \frac{\partial}{\partial z^a}, \\ \tilde{Z}^a &= \tilde{z}^a, & \tilde{W}^b &= (-)^{|a| \cdot |b|} \omega^{ba} \frac{\partial}{\partial \tilde{z}^a}. \end{aligned} \quad (5.42)$$

At quantum level, the classical constraint eqs(5.40) are replaced by operators acting on states $|\Psi_{h, \tilde{h}}\rangle$ of the Hilbert space of the fermionic twistor like system. More precisely,

we have,

$$\begin{aligned}\omega_{ab} \left(W^b Z^a + (-)^{|a| \cdot |b|} Z^a W^b \right) & \mid \Psi_{h, \tilde{h}} > = 4h \mid \Psi_{h, \tilde{h}} >, \\ \omega_{ab} \left(\widetilde{W}^b \widetilde{Z}^a + (-)^{|a| \cdot |b|} \widetilde{Z}^a \widetilde{W}^b \right) & \mid \Psi_{h, \tilde{h}} > = 4\tilde{h} \mid \Psi_{h, \tilde{h}} >,\end{aligned}\quad (5.43)$$

or equivalently by using $\Psi(z, \tilde{z}) = \langle z, \tilde{z} \mid \Psi \rangle$,

$$\begin{aligned}\omega_{ab} (-)^{|a| \cdot |b|} \left[\omega^{ba} + 2(-)^{|c| \cdot |b|} \omega^{bc} z^a \frac{\partial}{\partial z^c} \right] \Psi_{h, \tilde{h}}(z, \tilde{z}) &= 4h \Psi_{h, \tilde{h}}(z, \tilde{z}), \\ \omega_{ab} (-)^{|a| \cdot |b|} \left[\omega^{ba} + 2(-)^{|c| \cdot |b|} \omega^{bc} \tilde{z}^a \frac{\partial}{\partial \tilde{z}^c} \right] \Psi_{h, \tilde{h}}(z, \tilde{z}) &= 4\tilde{h} \Psi_{h, \tilde{h}}(z, \tilde{z}).\end{aligned}\quad (5.44)$$

The first term of this relation $\sum (-)^{|a| \cdot |b|} \omega_{ab} \omega^{ba}$ is proportional to the supertrace of 4×4 matrix,

$$Str(\omega_{ab} \omega^{ba}) = Str \begin{pmatrix} I & 0 \\ 0 & I \end{pmatrix}, \quad (5.45)$$

which vanishes identically. So one is left with the following,

$$\begin{aligned}\left(\lambda^\alpha \frac{\partial}{\partial \lambda^\alpha} + \xi_-^{\dot{\alpha}} \frac{\partial}{\partial \xi_-^{\dot{\alpha}}} \right) \Psi_{h, \tilde{h}}(z, \tilde{z}) &= 2h \Psi_{h, \tilde{h}}(z, \tilde{z}), \\ \left(\nu^{\dot{\alpha}} \frac{\partial}{\partial \nu^{\dot{\alpha}}} + \zeta_+^\alpha \frac{\partial}{\partial \zeta_+^\alpha} \right) \Psi_{h, \tilde{h}}(z, \tilde{z}) &= 2\tilde{h} \Psi_{h, \tilde{h}}(z, \tilde{z})\end{aligned}\quad (5.46)$$

showing that twistor space wave functions $\Psi_{h, \tilde{h}}(z, \tilde{z})$ are homogeneous superfunctions of degree $(2h, 2\tilde{h})$,

$$\Psi(e^{+\varphi} z, e^{+\theta} \tilde{z}) = e^{+(2h+2\tilde{h})\varphi} \Psi(z, \tilde{z}). \quad (5.47)$$

Charge quantization of scaling symmetry requires $2h$ and $2\tilde{h}$ to be integer,

$$2h \in \mathbb{Z}, \quad 2\tilde{h} \in \mathbb{Z}, \quad (5.48)$$

in agreement with local properties of quantum field theory requiring helicity to take half integer values. Moreover, wave function $\Psi_{h, \tilde{h}}(z, \tilde{z}) = \Psi_{h, \tilde{h}}(\lambda^\alpha, \nu^{\dot{\alpha}} \mid \zeta_+^\alpha, \xi_{-\dot{\alpha}})$ can be viewed as superfields with expansions type (2.67). We have

$$\begin{aligned}\Psi(z, \tilde{z}) &= \phi + \zeta_+^\alpha \psi_{-\alpha} + \xi_{-\dot{\alpha}} \psi_+^{\dot{\alpha}} + \zeta_+^2 F_{--} + \xi_-^2 F_{++} + \zeta_+^\alpha \xi_{-\dot{\alpha}} A_{\alpha}^{\dot{\alpha}} \\ &\quad + \xi_-^2 \zeta_+^\alpha \chi_{+\alpha} + \zeta_+^2 \xi_{-\dot{\alpha}} \chi_-^{\dot{\alpha}} + \zeta_+^2 \xi_-^2 \Delta\end{aligned}\quad (5.49)$$

where the component fields are as $\phi = \phi_{h, \tilde{h}}(\lambda^\alpha, \nu^{\dot{\alpha}})$ and so on. Using the transformation laws (5.19, 5.47), one can write down the corresponding ones for the component fields.

5.4 Comments

Here we want to make three comments regarding the study developed above:

(1) The analysis done for the $\mathbb{R}^{(2,2)}$ case with group $SO(2,2)$ can be naturally extended for Minkowski $\mathbb{R}^{(1,3)}$ and Euclidean $\mathbb{R}^4$ geometries with rotation groups $SO(1,3)$ and $SO(4)$ respectively. For the case of $\mathbb{R}^{(1,3)}$, spinors are in $SL(2, C)$ and the correspondence between generic superfield (5.49) and 4D $\mathcal{N} = 1$ supersymmetry representations is manifest.

(2) Target space supersymmetry is generated by the following superspace translations,

$$\begin{aligned}\delta\zeta_+^\alpha &\sim \varepsilon_+^\alpha, \\ \delta\xi_{-\dot{\alpha}} &\sim \varepsilon_{-\dot{\alpha}}, \\ \delta\lambda^\alpha &\sim \zeta_+^\alpha \varepsilon_{-\dot{\alpha}} \nu^{\dot{\alpha}} \\ \delta\nu^{\dot{\alpha}} &\sim \xi_{-\dot{\alpha}}^\alpha \lambda_\alpha\end{aligned}\tag{5.50}$$

where ε_+^α and $\varepsilon_{-\dot{\alpha}}$ are the target space supersymmetric parameters. These transformations are motivated by group covariance and dimensional arguments. Putting these naive transformations back into eq(5.49), we can get the supersymmetric transformations of the target space field variables. For the leading component fields, we have the following transformations,

$$\begin{aligned}\delta\phi &\sim \varepsilon_+^\alpha \psi_{-\alpha} + \varepsilon_{-\dot{\alpha}} \psi_+^{\dot{\alpha}} \\ \delta\psi_{-\alpha} &\sim 2\varepsilon_{+\alpha} F_{--} + \varepsilon_{-\dot{\alpha}} \left(\nu^{\dot{\alpha}} \frac{\partial}{\partial \lambda^\alpha} \phi - A_\alpha^{\dot{\alpha}} \right), \\ \delta\psi_+^{\dot{\alpha}} &\sim 2\varepsilon_{-\dot{\alpha}} F_{++} + \varepsilon_+^\alpha \left(\lambda_\alpha \frac{\partial}{\partial \nu_{\dot{\alpha}}} \phi + A_\alpha^{\dot{\alpha}} \right),\end{aligned}\tag{5.51}$$

and so on. Now performing a second supersymmetric transformation on $\delta\phi$ with supersymmetric parameters $\varepsilon'_{+\alpha}$ and $\varepsilon'_{-\dot{\alpha}}$, we have $\delta'\delta\phi \sim \varepsilon_+^\alpha \delta'\psi_{-\alpha} + \varepsilon_{-\dot{\alpha}} \delta'\psi_+^{\dot{\alpha}}$. Then replacing $\delta'\psi_{-\alpha}$ and $\delta'\psi_+^{\dot{\alpha}}$ by their expressions, we get

$$\begin{aligned}\delta'\delta\phi &\sim \left[2\varepsilon_+^\alpha \varepsilon'_{+\alpha} F_{--} + \varepsilon_+^\alpha \varepsilon'_{-\dot{\alpha}} \left(\nu^{\dot{\alpha}} \frac{\partial}{\partial \lambda^\alpha} \phi - A_\alpha^{\dot{\alpha}} \right) \right] \\ &\quad + \left[2\varepsilon_{-\dot{\alpha}} \varepsilon'^{\dot{\alpha}} F_{++} + \varepsilon_{-\dot{\alpha}} \varepsilon'_+{}^\alpha \left(\lambda_\alpha \frac{\partial}{\partial \nu_{\dot{\alpha}}} \phi + A_\alpha^{\dot{\alpha}} \right) \right]\end{aligned}\tag{5.52}$$

From this relation, one can compute the commutator $[\delta', \delta]\phi$, we find

$$\begin{aligned}[\delta', \delta]\phi &\sim (\varepsilon_+^\alpha \varepsilon'_{-\dot{\alpha}} - \varepsilon_+^{\alpha'} \varepsilon_{-\dot{\alpha}}) \nu^{\dot{\alpha}} \frac{\partial}{\partial \lambda^\alpha} \phi \\ &\quad + (\varepsilon_{-\dot{\alpha}} \varepsilon'^{\dot{\alpha}}_+ - \varepsilon'_{-\dot{\alpha}} \varepsilon_+^{\dot{\alpha}}) \lambda_\alpha \frac{\partial}{\partial \nu_{\dot{\alpha}}} \phi \\ &\quad + 2(\varepsilon_+^\alpha \varepsilon'_{+\alpha} - \varepsilon_+^{\alpha'} \varepsilon_{+\alpha}) F_{--} + 2(\varepsilon_{-\dot{\alpha}} \varepsilon'^{\dot{\alpha}}_- - \varepsilon'_{-\dot{\alpha}} \varepsilon_-^{\dot{\alpha}}) F_{++} \\ &\quad + [(\varepsilon_{-\dot{\alpha}} \varepsilon'^{\dot{\alpha}}_+ + \varepsilon_+^{\alpha'} \varepsilon_{-\dot{\alpha}}) - (\varepsilon'_{-\dot{\alpha}} \varepsilon_+^{\dot{\alpha}} + \varepsilon_+^\alpha \varepsilon'_{-\dot{\alpha}})] A_\alpha^{\dot{\alpha}}.\end{aligned}\tag{5.53}$$

But this relation may be simplified by using the following identities,

$$\begin{aligned}
\varepsilon_+^\alpha \varepsilon'_{+\alpha} &= \varepsilon_+^{\alpha'} \varepsilon_{+\alpha}, \\
\varepsilon_{-\dot{\alpha}} \varepsilon'^{\dot{\alpha}}_- &= \varepsilon'_{-\dot{\alpha}} \varepsilon_{-\dot{\alpha}}, \\
\varepsilon_{-\dot{\alpha}} \varepsilon'^{\dot{\alpha}}_+ &= -\varepsilon_+^{\alpha'} \varepsilon_{-\dot{\alpha}}, \\
\varepsilon'_{-\dot{\alpha}} \varepsilon_+^\alpha &= -\varepsilon_+^\alpha \varepsilon'_{-\dot{\alpha}} \\
(\varepsilon_+^\alpha \varepsilon'_{-\dot{\alpha}} - \varepsilon_+^{\alpha'} \varepsilon_{-\dot{\alpha}}) &= -(\varepsilon'_{-\dot{\alpha}} \varepsilon_+^\alpha - \varepsilon_{-\dot{\alpha}} \varepsilon'^{\dot{\alpha}}_+).
\end{aligned} \tag{5.54}$$

Taking into account these features, one ends with the result,

$$[\delta_{\varepsilon'}, \delta_\varepsilon] \phi \sim -(\varepsilon'_{-\dot{\alpha}} \varepsilon_+^\alpha - \varepsilon_{-\dot{\alpha}} \varepsilon'^{\dot{\alpha}}_+) P_\alpha^{\dot{\alpha}} \phi, \tag{5.55}$$

with

$$P_\alpha^{\dot{\alpha}} = \nu^{\dot{\alpha}} \frac{\partial}{\partial \lambda^\alpha} + \lambda_\alpha \frac{\partial}{\partial \nu_{\dot{\alpha}}}. \tag{5.56}$$

This supersymmetric transformation should hold also by the remaining component fields. As such target space supersymmetry can be defined in the twistor like space described above as follows,

$$\begin{aligned}
\{D_{-\alpha}, D_+^{\dot{\alpha}}\} &= 2 P_\alpha^{\dot{\alpha}}, \\
[P_\alpha^{\dot{\alpha}}, D_{-\alpha}] &= [P_\alpha^{\dot{\alpha}}, D_+^{\dot{\alpha}}] = 0.
\end{aligned} \tag{5.57}$$

Using the twistor like variables $(\lambda_\alpha, \nu_{\dot{\alpha}}, \zeta_+^\alpha, \xi_{-\dot{\alpha}})$, this superalgebra can be realized as,

$$\begin{aligned}
D_+^{\dot{\alpha}} &= \frac{\partial}{\partial \xi_{-\dot{\alpha}}} + \zeta_+^\alpha \lambda_\alpha \frac{\partial}{\partial \nu_{\dot{\alpha}}}, \\
D_{-\alpha} &= \frac{\partial}{\partial \zeta_+^\alpha} + \xi_{-\dot{\alpha}} \nu^{\dot{\alpha}} \frac{\partial}{\partial \lambda^\alpha}, \\
P_\alpha^{\dot{\alpha}} &= \nu^{\dot{\alpha}} \frac{\partial}{\partial \lambda^\alpha} + \lambda_\alpha \frac{\partial}{\partial \nu_{\dot{\alpha}}}.
\end{aligned} \tag{5.58}$$

Note that the above realization is not the unique one in twistor like space; we have in addition two more "chiral" ones. The first one corresponds to,

$$\begin{aligned}
D_+^{\dot{\alpha}} &= \frac{\partial}{\partial \xi_{-\dot{\alpha}}}, \\
D_{-\alpha} &= \frac{\partial}{\partial \zeta_+^\alpha} + 2 \xi_{-\dot{\alpha}} \nu^{\dot{\alpha}} \frac{\partial}{\partial \lambda^\alpha}, \\
P_\alpha^{\dot{\alpha}} &= \nu^{\dot{\alpha}} \frac{\partial}{\partial \lambda^\alpha}.
\end{aligned} \tag{5.59}$$

and the second one is given by the representation,

$$\begin{aligned}
D_+^{\dot{\alpha}} &= \frac{\partial}{\partial \xi_{-\dot{\alpha}}} + 2 \zeta_+^\alpha \lambda_\alpha \frac{\partial}{\partial \nu_{\dot{\alpha}}}, \\
D_{-\alpha} &= \frac{\partial}{\partial \zeta_+^\alpha}, \\
P_\alpha^{\dot{\alpha}} &= \lambda_\alpha \frac{\partial}{\partial \nu_{\dot{\alpha}}}.
\end{aligned} \tag{5.60}$$

Note also that in chiral eqs, the original energy momentum vector $P_\alpha^{\dot{\alpha}} = \left(\nu^{\dot{\alpha}} \frac{\partial}{\partial \lambda^\alpha} + \lambda_\alpha \frac{\partial}{\partial \nu_{\dot{\alpha}}} \right)$ of eqs(5.58) has been split into two parts. Each part appears in one of the two chiral realizations; i.e $\nu^{\dot{\alpha}} \frac{\partial}{\partial \lambda^\alpha}$ in (5.59) and $\lambda_\alpha \frac{\partial}{\partial \nu_{\dot{\alpha}}}$ in (5.60).

(3) The twistor like projective transformations (5.18) can be restricted by using the identity (5.21). In this case, the gauge fields $V(\tau)$ and $\tilde{V}_\tau(\tau)$ get identified and so the field action reduces to,

$$\begin{aligned} \mathcal{S}_{FT}[z, w, \tilde{z}, \tilde{w}, V] &= \frac{1}{2} \left(\int d\tau [w^b D_\tau z^a - (D_\tau w^b) z^a] \omega_{ab} \right) \\ &+ \frac{1}{2} \left(\int d\tau [\tilde{w}^b D_\tau \tilde{z}^a - (D_\tau \tilde{w}^b) \tilde{z}^a] \omega_{ab} \right) \\ &+ \int d\tau 2hV. \end{aligned} \quad (5.61)$$

By eliminating V through its equation of motion, we get the following constraint eq,

$$\omega_{ab} w^b z^a + \omega_{ab} \tilde{w}^b \tilde{z}^a - 2h = 0 \quad (5.62)$$

or equivalently by using (5.17),

$$(\nu_\alpha \lambda^\alpha - \zeta_{+\dot{\alpha}} \xi^{\dot{\alpha}}_-) - (\nu_{\dot{\alpha}} \lambda^\alpha - \zeta_{+\alpha} \xi^\alpha_-) - 2h = 0 \quad (5.63)$$

This constraint eq can be also rewritten as,

$$\omega_{ab} \left[\left(w^b z^a + (-)^{|a| \cdot |b|} z^a w^b \right) + \left(\tilde{w}^b \tilde{z}^a + (-)^{|a| \cdot |b|} \tilde{z}^a \tilde{w}^b \right) \right] - 4h = 0 \quad (5.64)$$

and at quantum level it gets promoted to,

$$\omega_{ab} \left[\left(W^b Z^a + (-)^{|a| \cdot |b|} Z^a W^b \right) + \tilde{W}^b \tilde{Z}^a + (-)^{|a| \cdot |b|} \tilde{Z}^a \tilde{W}^b \right] |\Psi\rangle = 4h |\Psi\rangle. \quad (5.65)$$

In differential representation $Z = z$, $W \sim \frac{\partial}{\partial z}$ and so on, this constraint reads as,

$$\omega_{ab} (-)^{|a| \cdot |b|} \left[(-)^{|c| \cdot |b|} \omega^{bc} z^a \frac{\partial}{\partial z^c} + (-)^{|c| \cdot |b|} \omega^{bc} \tilde{z}^a \frac{\partial}{\partial \tilde{z}^c} \right] \Psi(z, \tilde{z}) = 2h \Psi(z, \tilde{z}). \quad (5.66)$$

where we have used the identity $\sum (-)^{|a| \cdot |b|} \omega_{ab} \omega^{ba} = 0$. So one is left with the following,

$$\left(\lambda^\alpha \frac{\partial}{\partial \lambda^\alpha} + \nu^{\dot{\alpha}} \frac{\partial}{\partial \nu_{\dot{\alpha}}} + \zeta_+^\alpha \frac{\partial}{\partial \zeta_+^\alpha} + \xi_-^{\dot{\alpha}} \frac{\partial}{\partial \xi_-^{\dot{\alpha}}} \right) \Psi(z, \tilde{z}) = 2h \Psi(z, \tilde{z}) \quad (5.67)$$

showing that twistor space wave functions $\Psi(z, \tilde{z})$ are homogeneous superfunctions of degree $2h$, $\Psi(e^{+\varphi} z, e^{+\varphi} \tilde{z}) = e^{+2h\varphi} \Psi(z)$. Charge quantization requires $2h$ to be integer.

6 Conclusion and discussion

In this paper, we have developed a pure fermionic twistor like model capturing in a remarkable way the link between world line (world sheet) supersymmetry and target space one. This model describes a field theoretic system based on solving the nilpotency property

$$\Upsilon_{\pm}^2 = \epsilon_{\dot{\alpha}\dot{\beta}} \epsilon^{\alpha\beta} \Upsilon_{\pm\alpha}^{\dot{\alpha}} \Upsilon_{\pm\beta}^{\dot{\beta}} = 0, \quad \alpha, \dot{\alpha} = 1, 2, \quad (6.1)$$

of world line fermions type $\Upsilon_{\pm\alpha}^{\dot{\alpha}} = \Upsilon_{\pm\alpha}^{\dot{\alpha}}(\tau)$ and can be viewed as the analogue of the standard Penrose model for a massless particle moving in $4D$ space time $\mathbb{R}^{(d,4-d)}$ with $d = 2, 3, 4$.

The construction involves three main levels:

(1) Start with the world line field action $\mathcal{S}_F = \mathcal{S}_F[\Upsilon]$ of a fermionic particle parameterized by the world line fields $\Upsilon_{\pm\alpha}^{\dot{\alpha}}$ and $\tilde{\Upsilon}_{\pm\alpha}^{\dot{\alpha}}$. The latter fields ($\tilde{\Upsilon}_{\pm\alpha}^{\dot{\alpha}}$), which is predicted by the solving of the constraint eq(6.1), see also (6.5), have been supplemented in order to cover the full spectrum of 4D target space supersymmetry.

(2) Borrow techniques from Penrose formalism to solve eq(4.9) and then use the underlying free twistor like fields to build the fermionic twistor like model with action $\mathcal{S}_{FT} = \mathcal{S}_{FT}[\lambda, \zeta, \mu, \xi]$ dual to the previous $\mathcal{S}_F$ one. Here λ, ζ, μ and ξ stand respectively for the twistor like world fields $\lambda^{\alpha}(\tau)$, $\zeta_{\pm}^{\alpha}(\tau)$, $\mu^{\dot{\alpha}}(\tau)$ and $\xi_{\pm\dot{\alpha}}(\tau)$ together with their canonical conjugate. These fields were shown to be the projective coordinates of a supertwistor space whose explicit structure depends on the signature of target space time geometry $\mathbb{R}^{(d,4-d)}$.

(3) Quantize canonically the twistor like free fields and solve the underlying quantum gauge constraints to finish at the end of the analysis with superfields describing $4D$ $\mathcal{N} = 1$ target space supersymmetric representations. Fermionic twistor like models with $4D$ extended supersymmetry are also possible; they require implementation of more world fermions.

Among the basic results of this study we would like to mention the two following:

(i) Penrose construction dealing with massless particles $x^m = x^m(\tau)$ moving in $4D$ space-time $\mathbb{R}^{(d,4-d)}$ with energy momentum $p_m \sim p_{\dot{\alpha}}^{\alpha}$ turns out to be apply to more general systems. The point is that the usual Penrose incidence relation $\mu^{\dot{\alpha}} = ix_{\beta}^{\dot{\alpha}} \lambda^{\beta}$ eq(4.11) as well as the representation,

$$p_{\dot{\alpha}}^{\alpha} = \lambda^{\alpha} \lambda_{\dot{\alpha}}, \quad (6.2)$$

solving the condition $p^2 = 0$ have infinitely many analogues with the desired features. For instance, eq(6.2) is nothing but the leading term of a large class of world line fields belonging to the field family,

$$(\Pi_{(s_1, s_2)})_{\dot{\alpha}}^{\alpha} = (f_{s_1})^{\alpha} (g_{s_2})_{\dot{\alpha}}, \quad s = s_1 + s_2 = 0, \pm 1, \pm 2, \dots \quad (6.3)$$

The $\Pi_{(s_1, s_2)}$ fields with $s_1 s_2 \neq (2n+1)$ capture all the required properties of p_α^α needed for the building of twistor QFT. Like for p_α^α , the quadratic invariant $\Pi_{(s_1, s_2)}^2$ of these bispinors is given by

$$\Pi_{(s_1, s_2)}^2 = f_{s_1}^2 \cdot g_{s_2}^2, \quad (6.4)$$

and vanishes identically whenever at least one of the world line fields $f_{s_1} = f_{s_1}(\tau)$ or $g_{s_2} = g_{s_2}(\tau)$ is a boson. The next leading terms in the above family coming after p_α^α , and which is associated with the level $s_1 + s_2 = 0$, corresponds obviously to taking,

$$s_1 + s_2 = \pm 1. \quad (6.5)$$

Thus we have the four following,

$$\begin{aligned} (\Pi_{(\pm 1, 0)})_\alpha^\alpha &= \zeta_\pm^\alpha \tilde{\lambda}_{\dot{\alpha}} \sim \Upsilon_{\pm\alpha}^{\dot{\alpha}} \\ (\Pi_{(0, \pm 1)})_\alpha^\alpha &= \lambda^\alpha \tilde{\zeta}_{\pm\dot{\alpha}} \sim \tilde{\Upsilon}_{\pm\alpha}^{\dot{\alpha}}. \end{aligned} \quad (6.6)$$

Because of the identities $\lambda^\alpha \lambda_\alpha = 0$ and $\tilde{\lambda}_{\dot{\alpha}} \tilde{\lambda}^{\dot{\alpha}} = 0$, these world line fields obey the Fermi nilpotency property,

$$\text{Tr}(\Pi_{(\pm 1, 0)}^2) = 0, \quad \text{Tr}(\Pi_{(0, \pm 1)}^2) = 0, \quad (6.7)$$

capturing the known antisymmetry property of fermions wave function. In our concern, these relations are thought of as the analogue of the condition $p^2 = 0$ for massless particles in target space- time $\mathbb{R}^{(d, 4-d)}$, and so we can use Penrose method to deal with them.

(ii) Though the field action of the first level of the construction, (see also (4.34-4.39)),

$$\begin{aligned} \mathcal{S}_F \sim & \int_{\mathbb{R}} d\tau \left[\frac{1}{2} \left(\Upsilon_{-\alpha}^{\dot{\alpha}} \frac{\partial}{\partial \tau} \Upsilon_{+\beta}^{\dot{\beta}} - \Upsilon_{+\beta}^{\dot{\beta}} \frac{\partial}{\partial \tau} \Upsilon_{-\alpha}^{\dot{\alpha}} \right) \epsilon^{\alpha\beta} \epsilon_{\dot{\alpha}\dot{\beta}} \right] \\ & + \int_{\mathbb{R}} d\tau \left[\frac{1}{2} \left(\tilde{\Upsilon}_{-\alpha}^{\dot{\alpha}} \frac{\partial}{\partial \tau} \tilde{\Upsilon}_{+\beta}^{\dot{\beta}} - \tilde{\Upsilon}_{+\beta}^{\dot{\beta}} \frac{\partial}{\partial \tau} \tilde{\Upsilon}_{-\alpha}^{\dot{\alpha}} \right) \epsilon^{\alpha\beta} \epsilon_{\dot{\alpha}\dot{\beta}} \right], \end{aligned} \quad (6.8)$$

involves only world line fermions $\Upsilon_{\pm\alpha}^{\dot{\alpha}}$ and $\tilde{\Upsilon}_{\pm\alpha}^{\dot{\alpha}}$ (no world line bosons x_α^α nor its conjugate momentum p_α^α), its twistor like dual namely,

$$\begin{aligned} \mathcal{S}_{FT} \sim & \int d\tau \left[\frac{1}{2} (\nu_\alpha \partial_\tau \lambda^\alpha - \lambda^\alpha \partial_\tau \nu_\alpha) - \frac{1}{2} (\nu^{\dot{\alpha}} \partial_\tau \lambda_{\dot{\alpha}} - \lambda_{\dot{\alpha}} \partial_\tau \nu^{\dot{\alpha}}) \right] \\ & + \int d\tau \left[\frac{1}{2} (\xi_{-\alpha} \partial_\tau \zeta_+^\alpha + \zeta_+^\alpha \partial_\tau \xi_{-\alpha}) - \frac{1}{2} (\xi_{-\dot{\alpha}} \partial_\tau \zeta_{+\dot{\alpha}} + \zeta_{+\dot{\alpha}} \partial_\tau \xi_{-\dot{\alpha}}) \right]. \end{aligned} \quad (6.9)$$

involves however bosonic $(\lambda^\alpha, \nu_\alpha, \lambda_{\dot{\alpha}}, \nu^{\dot{\alpha}})$ and fermionic $(\zeta_+^\alpha, \xi_{-\alpha}, \zeta_{+\dot{\alpha}}, \xi_{-\dot{\alpha}})$ twistor like fields; thanks to the splitting (6.6). This feature, which shows that eq(6.8) is just a part of more general picture (see also last paragraph of this discussion); should be viewed as a novelty in twistor field theory. The reason is that starting with eq(6.8) and solve the

underlying constraint eqs, one discovers that the quantum spectrum of the dual fermionic twistor like model has automatically $4D$ target space supersymmetry.

In the end, we would like to note that the fermionic twistor construction developed in this paper depends on the signature of $4D$ target space $\mathbb{R}^{(d,4-d)}$ since corresponding spinors exhibit different properties. In the present study we have given explicit details for the case of particles moving in target space $\mathbb{R}^{(2,2)}$; the extension to the others geometries is straightforward. There, one also has the power of algebraic complex geometry. Note also that our analysis applies as well to world sheet fields and heterotic $(1,0)$ and $(0,1)$ QFT₂ models along the line described in sections 2 and 3. Note moreover that, though enough to recover $4D$ target space time supersymmetry, our pure fermionic model is in fact not completely independent from the standard Penrose bosonic one. Both of them may be viewed as just two parts of a world line supersymmetric theory. There, the fields $\Upsilon_{\pm\alpha}^{\dot{\alpha}}$ can be viewed as the fermionic partners of $x^m \sim x_{\alpha}^{\dot{\alpha}}$, that is interchanged under world line supersymmetry, $\delta_{susy} x_{\alpha}^{\dot{\alpha}} = \varepsilon_{\mp} \Upsilon_{\pm\alpha}^{\dot{\alpha}}$ and so on. This idea has been discussed shortly in present paper (sections 2 and 3). It would be interesting to deeper this issue for world line fields; but also for strings. Progress in this direction will be reported elsewhere.

Acknowledgement 2

This research work is supported by the program Protars III D12/25, CNRST. The authors thank ICTP (Senior Associate Programme and Organizers of Spring School) for kind hospitality and where part of this work has been done.

References

- [1] E. Witten, “Perturbative gauge theory as a string theory in twistor space”, Com. Math. Phys. 252 (2004) 189, hep-th/0312171.
- [2] E. Witten, “Parity invariance for strings in twistor space”, hep-th/0403199.
- [3] I. Bars, “Twistor superstring in 2T-physics,” Phys. Rev. D70 (2004) 104022, hep-th/0407239.
- [4] Freddy Cachazo, Peter Svrcek, Lectures on Twistor Strings and Perturbative Yang-Mills Theory, PoS RTN2005 (2005) 004, hep-th/0504194
- [5] I. Bena, Z. Bern and D. A. Kosower, “Twistor-space recursive formulation of gauge theory amplitudes,” Phys. Rev. D 71, 045008 (2005), hep-th/0406133.

- [6] S. J. Parke and T. R. Taylor, “Perturbative QCD utilizing extended supersymmetry,” *Phys. Lett.* B157 (1985) 81.
- [7] S. J. Parke and T. R. Taylor, “An amplitude for n gluon scattering,” *Phys. Rev. Lett.* 56 (1986) 2459.
- [8] Z. Bern, L. J. Dixon, and D. A. Kosower, “One loop corrections to five gluon amplitudes,” *Phys. Rev. Lett.* 70 (1993) 2677–2680, hep-ph/9302280.
- [9] Z. Bern, A. De Freitas, and L. J. Dixon, “Two-loop helicity amplitudes for gluon gluon scattering in QCD and supersymmetric Yang-Mills theory,” *JHEP* 03 (2002) 018, hep-ph/0201161.
- [10] Z. Bern, A. De Freitas, and L. J. Dixon, “Two-loop helicity amplitudes for quark gluon scattering in QCD and gluino gluon scattering in supersymmetric Yang-Mills theory,” *JHEP* 06 (2003) 028, hep-ph/0304168.
- [11] N. Berkovits and E. Witten, Conformal Supergravity in Twistor-String Theory, *JHEP* 0408 (2004) 009, hep-th/0406051.
- [12] N. Berkovits, “An Alternative string theory in twistor space for $N=4$ Super Yang-Mills”, *Phys. Rev. Lett.* 93 (2004) 011601, hep-th/0402045.
- [13] N. Berkovits and L. Motl, “Cubic twistorial string field theory”, *JHEP* 0404 (2004) 56, hep-th/0403187.
- [14] L.J.Mason, Twistor actions for non-self-dual fields; a new foundation for twistor-string theory, *JHEP* 0510 (2005) 009, hep-th/0507269
- [15] Christian Saemann, Aspects of Twistor Geometry and Supersymmetric Field Theories within Superstring Theory, hep-th/0603098,
- [16] R. Penrose, “Twistor Algebra,” *J. Math. Phys.* 8 (1967) 345; “The Twistor Program,” *Rept. Math. Phys.* 12 (1977) 65.
- [17] R. Penrose and M.A. MacCallum, “An approach to the quantization of fields and space-time”, *Phys. Rept.* C6 (1972) 241;
- [18] R. Penrose and W. Rindler, *Spinors and space-time II*, Cambridge Univ. Press (1986).
- [19] R. Ahl Laamara, A. Belhaj, L.B. Drissi, E.H. Saidi, On Local Calabi-Yau Supermanifolds and Their Mirrors, *J.Phys.* A39 (2006) 5965-5978, hep-th/0601215

- [20] A. Belhaj, L.B. Drissi, J. Rasmussen, E.H. Saidi, A. Sebbar, Toric Calabi-Yau supermanifolds and mirror symmetry, J.Phys. A38 (2005) 6405-6418, hep-th/0410291
- [21] Shigenori Seki, Katsuyuki Sugiyama, Tatsuya Tokunaga, Superconformal Symmetry in Linear Sigma Model on Supermanifolds, hep-th/0605021.
- [22] Olaf Lechtenfeld, Christian Saemann, Matrix Models and D-branes in Twistor String Theory, JHEP 0603 (2006) 002, hep-th/0511130,
- [23] Alexander D. Popov, Christian Saemann, On Supertwistors, the Penrose-Ward Transform and $\mathcal{N} = 4$ super Yang-Mills Theory, Adv.Theor.Math.Phys. 9 (2005) 931-998, hep-th/0405123,
- [24] Pierre Cartier, Cecile DeWitt-Morette, Matthias Ihl, Christian Saemann, Maria E. Bell, Supermanifolds - Application to Supersymmetry, math-ph/0202026, to appear in the "Michael Marinov Memorial Volume", "Multiple facets of quantization and supersymmetry", Eds. M. Olshanetsky and A. Vainshtein, World Scientific, 2002:
- [25] Martin Wolf, On hidden symmetries of a super gauge theory and twistor string theory, JHEP 0502:018,2005, hep-th/0412163.
- [26] Riccardo Ricci, Super Calabi-Yau's and special Lagrangians, YITP-SB-05-42, hep-th/0511284
- [27] Houda Jehjouh, Memoire de DESA-PHE, Lab/UFR-High Energy Physics Fac Sciences, Rabat, september (2005),
- [28] From twistor string theory to $4D \mathcal{N} = 4$ conformal supergravity, to appear in African Journal of Mathematical Physics AJMP (2006)
- [29] Andrew Neitzke, Cumrun Vafa, $\mathcal{N} = 2$ strings and the twistorial Calabi-Yau, hep-th/0402128,
- [30] Robbert Dijkgraaf, Sergei Gukov, Andrew Neitzke, Cumrun Vafa, Topological M-theory as Unification of Form Theories of Gravity, hep-th/0411073
- [31] El Hassan Saidi, Moulay Brahim Sedra, Topological string in harmonic space and correlation functions in S^3 stringy cosmology, hep-th/0604204, to appear in Nuclear Physics B.
- [32] Alexander D. Popov, Christian Saemann, Martin Wolf, The Topological B-model on a Mini-Supertwistor Space and Supersymmetric Bogomolny Monopole Equations, JHEP 0510 (2005) 058, hep-th/0505161

- [33] Rachid Ahl Laamara, Lalla Btissam Drissi, El Hassan Saidi, D-string fluid in conifold: I. Topological gauge model, Nucl Phys B743 (2006) 333, hep-th/0604001
- [34] El Hassan Saidi, Topological $SL(2)$ Gauge Theory on Conifold, hep-th/0601020
- [35] Rachid Ahl Laamara, Lalla Btissam Drissi, El Hassan Saidi, D-string fluid in conifold: II. Matrix model proposal for droplets on S^3 and S^2 , Lab/UFR-HEP/06/01, GNPHE/06/01, VACBT/06/01.
- [36] I. Bars, “Twistors and 2T-physics,” AIP Conf. Proc. 767 (2005) 3, hep-th/0502065
- [37] Itzhak Bars, Moises Picon, Twistor Transform in d Dimensions and a Unifying Role for Twistors, Phys.Rev. D73 (2006) 064033, hep-th/0512348